



The University of
Nottingham

UNITED KINGDOM • CHINA • MALAYSIA

Debnath, Ujjwal Kanti (2010) Factors predicting the outcome following treatment for lumbar spondylolysis. DM thesis, University of Nottingham.

Access from the University of Nottingham repository:

http://eprints.nottingham.ac.uk/12780/5/Levels_of_evidence.pdf

Copyright and reuse:

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

This article is made available under the University of Nottingham End User licence and may be reused according to the conditions of the licence. For more details see:
http://eprints.nottingham.ac.uk/end_user_agreement.pdf

For more information, please contact eprints@nottingham.ac.uk

Oxford Centre for Evidence-based Medicine Levels of Evidence (May 2001)

Level	Therapy/Prevention, Aetiology/Harm	Prognosis	Diagnosis	Differential diagnosis/symptom prevalence study	Economic and decision analyses
1a	SR (with <u>homogeneity*</u>) of RCTs	SR (with <u>homogeneity*</u>) of inception cohort studies; <u>CDR†</u> validated in different populations	SR (with homogeneity*) of Level 1 diagnostic studies; <u>CDR†</u> with 1b studies from different clinical centres	SR (with homogeneity*) of prospective cohort studies	SR (with homogeneity*) of Level 1 economic studies
1b	Individual RCT (with narrow <u>Confidence Interval‡</u>)	Individual inception cohort study with ≥ 80% follow-up; <u>CDR†</u> validated in a single population	Validating** cohort study with good††† reference standards; or <u>CDR†</u> tested within one clinical centre	Prospective cohort study with good follow-up****	Analysis based on clinically sensible costs or alternatives; systematic review(s) of the evidence; and including multi-way sensitivity analyses
1c	<u>All or none§</u>	All or none case-series	Absolute SpPins and SnNouts††	All or none case-series	Absolute better-value or worse-value analyses ††††
2a	SR (with <u>homogeneity*</u>) of cohort studies	SR (with <u>homogeneity*</u>) of either retrospective cohort studies or untreated control groups in RCTs	SR (with homogeneity*) of Level >2 diagnostic studies	SR (with homogeneity*) of 2b and better studies	SR (with homogeneity*) of Level >2 economic studies
2b	Individual cohort study (including low quality RCT; e.g., <80% follow-up)	Retrospective cohort study or follow-up of untreated control patients in an RCT; Derivation of <u>CDR†</u> or validated on split-sample§§§ only	Exploratory** cohort study with good††† reference standards; <u>CDR†</u> after derivation, or validated only on split-sample§§§ or databases	Retrospective cohort study, or poor follow-up	Analysis based on clinically sensible costs or alternatives; limited review(s) of the evidence, or single studies; and including multi-way sensitivity analyses
2c	"Outcomes" Research; Ecological studies	"Outcomes" Research		Ecological studies	Audit or outcomes research
3a	SR (with <u>homogeneity*</u>) of case-control studies		SR (with homogeneity*) of 3b and better studies	SR (with homogeneity*) of 3b and better studies	SR (with homogeneity*) of 3b and better studies
3b	Individual Case-Control Study		Non-consecutive study; or without consistently applied reference standards	Non-consecutive cohort study, or very limited population	Analysis based on limited alternatives or costs, poor quality estimates of data, but including sensitivity analyses incorporating clinically sensible variations.
4	Case-series (and <u>poor quality cohort and case-control studies§§</u>)	Case-series (and <u>poor quality prognostic cohort studies***</u>)	Case-control study, poor or non-independent reference standard	Case-series or superseded reference standards	Analysis with no sensitivity analysis
5	Expert opinion without explicit critical appraisal, or based on physiology, bench research or "first principles"	Expert opinion without explicit critical appraisal, or based on physiology, bench research or "first principles"	Expert opinion without explicit critical appraisal, or based on physiology, bench research or "first principles"	Expert opinion without explicit critical appraisal, or based on physiology, bench research or "first principles"	Expert opinion without explicit critical appraisal, or based on economic theory or "first principles"

Produced by Bob Phillips, Chris Ball, Dave Sackett, Doug Badenoch, Sharon Straus, Brian Haynes, Martin Dawes since November 1998.

Notes

Users can add a minus-sign "-" to denote the level of that fails to provide a conclusive answer because of:

- EITHER a single result with a wide Confidence Interval (such that, for example, an ARR in an RCT is not statistically significant but whose confidence intervals fail to exclude clinically important benefit or harm)
- OR a Systematic Review with troublesome (and statistically significant) heterogeneity.
- Such evidence is inconclusive, and therefore can only generate Grade D recommendations.

*	By homogeneity we mean a systematic review that is free of worrisome variations (heterogeneity) in the directions and degrees of results between individual studies. Not all systematic reviews with statistically significant heterogeneity need be worrisome, and not all worrisome heterogeneity need be statistically significant. As noted above, studies displaying worrisome heterogeneity should be tagged with a "-" at the end of their designated level.
†	Clinical Decision Rule. (These are algorithms or scoring systems which lead to a prognostic estimation or a diagnostic category.)
‡	See note #2 for advice on how to understand, rate and use trials or other studies with wide confidence intervals.
§	Met when <u>all</u> patients died before the Rx became available, but some now survive on it; or when some patients died before the Rx became available, but <u>none</u> now die on it.
§§	By poor quality <u>cohort</u> study we mean one that failed to clearly define comparison groups and/or failed to measure exposures and outcomes in the same (preferably blinded), objective way in both exposed and non-exposed individuals and/or failed to identify or appropriately control known confounders and/or failed to carry out a sufficiently long and complete follow-up of patients. By poor quality <u>case-control</u> study we mean one that failed to clearly define comparison groups and/or failed to measure exposures and outcomes in the same (preferably blinded), objective way in both cases and controls and/or failed to identify or appropriately control known confounders.
§§§	Split-sample validation is achieved by collecting all the information in a single tranche, then artificially dividing this into "derivation" and "validation" samples.
††	An "Absolute SpPin" is a diagnostic finding whose <u>Specificity</u> is so high that a <u>Positive</u> result rules-in the diagnosis. An "Absolute SnNout" is a diagnostic finding whose <u>Sensitivity</u> is so high that a <u>Negative</u> result rules-out the diagnosis.
‡‡	Good, better, bad and worse refer to the comparisons between treatments in terms of their clinical risks and benefits.
†††	<u>Good</u> reference standards are independent of the test, and applied blindly or objectively to applied to all patients. <u>Poor</u> reference standards are haphazardly applied, but still independent of the test. Use of a non-independent reference standard (where the 'test' is included in the 'reference', or where the 'testing' affects the 'reference') implies a level 4 study.
††††	Better-value treatments are clearly as good but cheaper, or better at the same or reduced cost. Worse-value treatments are as good and more expensive, or worse and the equally or more expensive.
**	Validating studies test the quality of a specific diagnostic test, based on prior evidence. An exploratory study collects information and trawls the data (e.g. using a regression analysis) to find which factors are 'significant'.
***	By poor quality prognostic cohort study we mean one in which sampling was biased in favour of patients who already had the target outcome, or the measurement of outcomes was accomplished in <80% of study patients, or outcomes were determined in an unblinded, non-objective way, or there was no correction for confounding factors.
****	Good follow-up in a differential diagnosis study is >80%, with adequate time for alternative diagnoses to emerge (eg 1-6 months acute, 1 - 5 years chronic)

Grades of Recommendation

A	consistent level 1 studies
B	consistent level 2 or 3 studies or extrapolations from level 1 studies
C	level 4 studies or extrapolations from level 2 or 3 studies
D	level 5 evidence or troublingly inconsistent or inconclusive studies of any level

"Extrapolations" are where data is used in a situation which has potentially clinically important differences than the original study situation.