



The University of
Nottingham

UNITED KINGDOM • CHINA • MALAYSIA

Wise, A.E. and Tarlinton, R.E. (2012) Seroprevalence of vectorborne diseases in free-roaming dogs in Goa, India. *Veterinary Record*, 170 (3). ISSN 0042-4900

Access from the University of Nottingham repository:

<http://eprints.nottingham.ac.uk/2472/1/Tarlintongoa.pdf>

Copyright and reuse:

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

This article is made available under the Creative Commons Attribution Non-commercial licence and may be reused according to the conditions of the licence. For more details see: <http://creativecommons.org/licenses/by-nc/2.5/>

A note on versions:

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact eprints@nottingham.ac.uk

Short Communications

Seroprevalence of vectorborne diseases in free-roaming dogs in Goa, India

A. E. Wise, R. E. Tarlinton

A majority of the Indian dogs are 'community dogs', semiferal dogs which are partially dependent on the human population and feral dogs. Communicable diseases may be a problem due to overcrowding, minimal veterinary care, and a climate favouring parasites and vectors. Some reports suggest that the vectorborne diseases, such as filariasis, babesiosis and ehrlichiosis, are endemic throughout India. However, there is little information concerning disease epidemiology in Indian dogs (Megat Abd Rani and others 2010a).

Vectorborne diseases of dogs are of particular interest, as pathogens such as *Borrelia burgdorferi* sensu lato (the agent of Lyme disease) are a zoonotic risk in addition to causing disease in their canine hosts (Megat Abd Rani and others 2010a). The pathogens examined here, the tickborne *B burgdorferi* sensu lato, *Ehrlichia canis*, *Anaplasma platys* and the mosquito-borne *Dirofilaria immitis*, are regarded as endemic in the dog populations in warm climate zones, worldwide.

The reported prevalence of *E canis* within India varies with 0.35 per cent and 18.9 per cent of dogs identified with canine ehrlichiosis by stained blood smears in Punjab and Nagpur, respectively (Juyal and others 1994, Samaradni and others 2003, Megat Abd Rani and others 2010a). Studies in Chennai reported that 50 per cent of privately-owned dogs tested positive for *E canis* when using species-specific PCR compared with 19 per cent by microscopy (Lakshmanan and others 2007). Megat Abd Rani and others (2011) reported a PCR-based prevalence of 27.2 to 39.5 per cent of *E canis* in tropical and subtropical Delhi and Mumbai, but an absence of this pathogen in the more temperate climate zones of north-West Bengal and Jammu Kashmir.

D immitis has been reported in northern India. Borthakur and others (2006) identified 34 per cent of 240 dogs at a slaughterhouse in north-east India to be infected with *D immitis*, and Megat Abd Rani and others (2010b) reported 4.3 per cent of dogs in Delhi to be positive by PCR-based tests. Indian veterinarians believe *D immitis* to be confined to north-east India. However, potential vectors, such as *Aedes albopictus* (the Asian tiger mosquito), for *D immitis*, are present throughout India (Megat Abd Rani and others 2010b).

There has been at least one study that has failed to identify *B burgdorferi* sensu lato in India (Handa and others 1999). *A platys* has recently been identified in Indian dogs by PCR at prevalences of 8 to




FIG 1: Seroprevalence of Dogs to *Dirofilaria immitis*, *Anaplasma platys*, *Ehrlichia canis* and *Borrelia burgdorferi* (N=48)

13 per cent in Mumbai and Delhi but is absent in more temperate climate zones (Megat Abd Rani and others 2011).

Blood samples from 48 dogs undergoing surgical sterilisation as part of an animal birth control programme at Animal Tracks, a Veterinary Centre run by the International Animal Rescue in North Goa, India, were tested for *D immitis* antigen, *B burgdorferi* sensu lato, *Anaplasma* species and *E canis* antibody using the SNAP 4Dx Test (IDEXX) kit according to the manufacturer's instructions. Biometric data including age, weight, sex and body condition on a five-point scale, and the area from which the dog was captured, were recorded for each animal. A summary of the samples collected is shown in Table 1. Samples were collected over an eight-week period in August and September 2011.

This project was approved by the University of Nottingham, School of Veterinary Medicine and Science, non-ASP (animals (scientific procedures) act) ethical committee.

Statistical testing of the associations between the biometric parameters and the disease was performed by chi-squared analysis using the Minitab version 15.1.0.1 (Minitab) statistical package.

Of the 48 dogs tested, 21 per cent (10) and 19 per cent (nine) tested positive for *A platys* and *E canis*, respectively. Co-infection with these two species was found in 10 per cent (five of 48) of the dogs tested. No cases of *D immitis* or *B burgdorferi* were found (Fig 1). There was a significant association for co-infection with *A platys* and *E canis* ($P < 0.005$). No other significant associations were found.

This short communication provides further evidence for the presence of *A platys* in Indian dogs with a seroprevalence of 21 per cent. This is very similar to the recent report by Megat Abd Rani and others (2011) of a PCR-based prevalence of 27.2 per cent for *A platys* in Mumbai, which is only 600 km north of Goa and in a similar climate zone. The prevalence of *E canis* (19 per cent) in this short communication is consistent with the other studies on this pathogen in India, which suggest prevalences from 0.35 to 50 per cent (Juyal and others 1994, Samaradni and others 2003, Lakshmanan and others 2007, Megat Abd Rani and others 2010a, Megat Abd Rani and others 2011).

Co-infection of *A platys* and *E canis* was found in 10 per cent of dogs; this association was statistically significant ($P < 0.005$). This would suggest transmission by a common vector, most likely *Rhipicephalus sanguineus*, which is known to be the vector of *E canis* (Nicholson and others 2010), and thought to be the vector of *A platys* (Yabsley and others 2008). *Rhipicephalus* species ticks are also known to form almost 100 per cent of the tick infestations in street dogs in the urban areas of India (Megat Abd Rani and others 2011) with up to 80 per cent of the dogs infested. The study also reported a similar rate of co-infections with *A platys* and *E canis* (4.5 to 7 per cent) in Mumbai and Delhi, respectively. The IDEXX 4Dx kit used in the present study is unable to distinguish between *A platys* and *Anaplasma*

Veterinary Record (2012) 170, 76a

doi: 10.1136/vr.100169

A. E. Wise,
R. E. Tarlinton, BVSc, PhD, MRCVS,
School of Veterinary Medicine and
Science, University of Nottingham,
Sutton Bonington Campus,
Loughborough, Leicestershire LE12
5PE, UK

Correspondence to Dr Tarlinton, e-mail:
rachel.tarlinton@nottingham.ac.uk

Provenance: not commissioned;
externally peer reviewed

Accepted September 27, 2011

Published Online First October 29, 2011



Short Communications

TABLE 1: Summary of the morphometric data of dogs sampled for this study

Number of animals	Mean (se) body condition score	Mean (se) bodyweight (kg)	% owned	
Adults (>2 years)				
Male	13	2.5 (0.1)	12.7 (0.6)	10
Female	9	2.8 (0.2)	14.4 (0.8)	20
Juveniles (≤2 years)				
Male	5	2 (0.0)	11.6 (0.7)	20
Female	21	2.4 (0.1)	12.6 (0.7)	36
Total	48			

phagocytophilum; however, given that Megat Abd Rani and others (2011) identified dogs infected with *A. platys* by PCR in a nearby geographical location at a similar prevalence, the seroreaction in the present study is likely to be due to this species and the authors have assumed this throughout this short communication.

No cases of *B. burgdorferi* were found in the dogs in this study. This may be a factor of the small number (48) of animals tested; however, there have been other studies in India reporting the absence of this pathogen (Handa and others 1999). Either the test kits are not able to detect Indian strains of this pathogen or it may be genuinely absent from India despite the presence of suitable vectors. Resolution of this question would require direct testing for the pathogen, either by culture or PCR rather than the indirect serological method utilised here.

One false-positive test for *D. immitis* was recorded in this study. Microfilaria were observed by light microscopy in thick blood smears from one dog that tested negative for *D. immitis* (data not shown); these were assumed to be *Dirofilaria repens* or *Acanthocheilonema reconditum*, species that have been reported in southern India (Ananda and others 2002, Sabu and others 2005, Megat Abd Rani and others 2010b). Crossreactions with *D. repens* have been reported in ELISA tests (Schrey and Trautvetter 1998). *D. immitis* is thought to be confined to North India (Borthakur and others 2006, Megat Abd Rani and others 2010b). The present study supports this theory.

This study has provided evidence for the presence of *A. platys* and *E. canis* in community dogs in Goa that will allow local veterinarians to presumptively treat dogs showing clinical signs of ehrlichiosis or anaplasmosis. The high prevalences of these parasites would indicate a significant risk for tickborne diseases in both the human and dog populations in this area, though the exact vector and pathogen systems involved require further work to be clarified.

Acknowledgements

The authors thank Carl Regan for his help and support in data collection and throughout the project; Astrid Almeida, all the staff and volunteers at Animal Tracks veterinary centre for their help and enthusiasm for this project; and the Universities Federation for Animal Welfare, University of Nottingham and the University Association Fund Award for their vital funding.

References

- ANANDA, K. J., D'SOUZAPLACID, E. & JAGANNATH, M. S. (2002) Methods for identification of microfilaria of *Dirofilaria repens* and *Dipetalonema reconditum*. *Veterinary Parasitology* **20**, 45-47
- BORTHAKUR, S. K., SARMAH, K., RAJKHOWA, T. K., DAS, M. R. & RAHMAN, S. (2006) *Dirofilaria immitis* infection in dog. *Journal of Veterinary Parasitology* **20**, 167-169
- HANDA, R., WALL, J. P., SINGH, S. & AGGARWAL, P. (1999) A prospective study of Lyme arthritis in north India. *Indian Journal of Medical Research* **110**, 107-109
- JUYAL, P. D., KALRA, I. S. & SINGHLA, L. D. (1994) Prevalence of haemoprotozoans in domestic animals in Punjab. In 6th National Congress of Veterinary Parasitology
- LAKSHMANAN, B., JOHN, L., GOMATHINAYAGAM, S. & DHINAKARRAJ, G. (2007) Molecular detection of *Ehrlichia canis* from blood of naturally infected dogs in India. *Veterinarski Arhiv* **77**, 307-312
- MEGAT ABD RANI, P. A., IRWIN, P. J., GATNE, M., COLEMAN, G. T., MCINNES, L. M. & TRAUB, R. J. (2010a) Canine vector borne diseases in India: a review of the literature and identification of existing knowledge gaps. *Parasites and Vectors* **3**, 28
- MEGAT ABD RANI, P. A., IRWIN, P. J., GATNE, M., COLEMAN, G. T., MCINNES, L. M. & TRAUB, R. J. (2010b) A survey of canine filarial diseases of veterinary and public health significance in India. *Parasites & Vectors* **3**, 30
- MEGAT ABD RANI, P. A., IRWIN, P. J., COLEMAN, G. T., GATNE, M. & TRAUB, R. J. (2011) A survey of canine tick borne diseases in India. *Parasites and Vectors* **4**, 141
- NICHOLSON, W. L., ALLEN, K. E., MCQUISTON, J. H., BREITSCHWERDT, E. B. & LITTLE, S. E. (2010) The increasing recognition of rickettsial pathogens in dogs and people. *Trends in Parasitology* **26**, 205-212
- SABU, L., DEVADA, K. & SUBRAMANIAN, H. (2005) Dirofilariosis in dogs and humans in Kerala. *Indian Journal of Medical Research* **121**, 691-693
- SAMARADNI, D., MASKE, D. K., KOLTE, S. W. & SHINDE, P. N. (2003) Ehrlichiosis in dogs in Nagpur. *Veterinary Parasitology* **17**, 165-166
- SCHREY, C. F. & TRAUTVETTER, E. (1998) Canine and feline heartworm disease—diagnosis and therapy. *Waltham Focus* **8**, 23-31
- YABSLEY, M. J., MCKIBBEN, J., MACPHERSON, C. N., CATTAN, P. E., CHERRY, N. A., HEGARTY, B. C., BREITSCHWERDT, E. B., O'CONNOR, T., & OTHERS (2008) Prevalence of *Ehrlichia canis*, *Anaplasma platys*, *Babesia canis vogeli*, *Hepatozoon canis*, *Bartonella vinsonii berkhoffii*, and *Rickettsia* spp. in dogs from Grenada. *Veterinary parasitology* **151**, 279-285

Seroprevalence of vectorborne diseases in free-roaming dogs in Goa, India

A. E. Wise and R. E. Tarlinton

Veterinary Record 2012 170: 76 originally published online October 29, 2011

doi: 10.1136/vr.100169

Updated information and services can be found at:

<http://veterinaryrecord.bmj.com/content/170/3/76.1.full.html>

These include:

- | | |
|-------------------------------|---|
| References | This article cites 12 articles
http://veterinaryrecord.bmj.com/content/170/3/76.1.full.html#ref-list-1 |
| Open Access | This paper is freely available online under the BMJ Journals unlocked scheme, see http://veterinaryrecord.bmj.com/info/unlocked.dtl |
| Email alerting service | Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article. |

Topic Collections

Articles on similar topics can be found in the following collections

[Open access](#) (39 articles)

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>