

White, Kate L. and Paine, Stuart and Harris, John (2017)
A clinical evaluation of the pharmacokinetics and
pharmacodynamics of intravenous alfaxalone in
cyclodextrin in male and female rats following a loading
dose and constant rate infusion. *Veterinary Anaesthesia
and Analgesia*, 44 (4). pp. 865-875. ISSN 1467-2987

Access from the University of Nottingham repository:

<http://eprints.nottingham.ac.uk/40482/1/1-s2.0-S1467298717300624-main.pdf>

Copyright and reuse:

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

This article is made available under the University of Nottingham End User licence and may be reused according to the conditions of the licence. For more details see:

http://eprints.nottingham.ac.uk/end_user_agreement.pdf

A note on versions:

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact eprints@nottingham.ac.uk

Accepted Manuscript

A clinical evaluation of the pharmacokinetics and pharmacodynamics of intravenous alfaxalone in cyclodextrin in male and female rats following a loading dose and constant rate infusion


Kate L. White, Stuart Paine, John Harris

PII: S1467-2987(17)30062-4

DOI: [10.1016/j.vaa.2017.01.001](https://doi.org/10.1016/j.vaa.2017.01.001)

Reference: VAA 88

To appear in: *Veterinary Anaesthesia and Analgesia*



Please cite this article as: White KL, Paine S, Harris J, A clinical evaluation of the pharmacokinetics and pharmacodynamics of intravenous alfaxalone in cyclodextrin in male and female rats following a loading dose and constant rate infusion, *Veterinary Anaesthesia and Analgesia* (2017), doi: 10.1016/j.vaa.2017.01.001.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 RESEARCH PAPER

2 **A clinical evaluation of the pharmacokinetics and pharmacodynamics of**
3 **intravenous alfaxalone in cyclodextrin in male and female rats following a loading**
4 **dose and constant rate infusion**

5 Kate L White*, Stuart Paine* & John Harris†

6

7 *School of Veterinary Medicine and Science, University of Nottingham, Sutton
8 Bonington, UK

9 †School of Biosciences, University of Nottingham, Sutton Bonington, UK

10

11 **Correspondence:** Kate White, School of Veterinary Medicine and Science, University
12 of Nottingham, Sutton Bonington, Loughborough, Leics LE12 5RD, UK. E-mail:
13 Kate.White@nottingham.ac.uk

14

15 *Running head: Alfaxalone anaesthesia in male and female rats*

16

17 **Abstract**

18 **Objective** To characterise, as a clinical study, the pharmacokinetics and
19 pharmacodynamics and describe the hypnotic effect of the neurosteroid alfaxalone (3 α -
20 hydroxy-5 α -pregnane-11, 20-dione) formulated with 2-hydroxypropyl- β -cyclodextrin
21 in male and female rats.

22 **Study design** Prospective, experimental laboratory study.

23 **Animals** Twelve (six male and six female) adult, aged matched Sprague Dawley rats.

24 **Methods** Surgery and instrumentation was performed under isoflurane anaesthesia in an
25 oxygen/nitrous oxide mixture (1:2) and local anaesthetic infiltration. All animals
26 received a loading dose (1.67 mg kg⁻¹ minute⁻¹) for 2.5 minutes followed by a constant
27 rate infusion (0.75 mg kg⁻¹ minute⁻¹) for 120 minutes of alfaxalone. Isoflurane and
28 nitrous oxide was discontinued 2.5 minutes after the alfaxalone infusion started.
29 Cardiorespiratory variables (heart rate, respiratory rate, arterial blood pressure, end tidal
30 carbon dioxide tension) and clinical signs of anaesthetic depth were evaluated
31 throughout anaesthesia. Carotid artery blood samples were collected at strategic time
32 points for blood gas analysis, haematology and biochemistry and plasma concentrations
33 of alfaxalone. Plasma samples were assayed using liquid chromatography-mass
34 spectrometry (LC/MS).

35 **Results** There were significant differences between the sexes for plasma clearance ($p =$
36 0.0008), half-life ($p = 0.0268$) and mean residence time ($p = 0.027$). Mean arterial blood
37 pressure was significantly higher in the male rats ($p = 0.0255$).

38 **Conclusions and clinical relevance** This study confirms alfaxalone solubilized in a 2-
39 hydroxypropyl- β -cyclodextrin provides excellent total intravenous anaesthesia in rats.

40 Sex-based differences in pharmacokinetics and pharmacodynamics were demonstrated
41 and must be considered when designing biomedical research models using alfaxalone.

42

43 *Keywords* alfaxalone, anaesthetics, intravenous, rat, steroid,

44

Introduction

Alfaxalone is a neuroactive steroid that modulates neurotransmission through interaction with a steroid recognition site on the GABA_A receptor complex causing a positive allosteric modulation of the ligand gated chloride channel resulting in inhibition of neuronal excitability (Harrison & Simmonds 1984; Turner et al. 1989). Such agents therefore have roles in anaesthesia, epilepsy, anxiety, insomnia, migraine and drug dependence (Rupprecht & Holsboer 1999). Alfaxalone had been used as an anaesthetic induction agent in humans and veterinary species for almost half a century but anaphylactoid reactions attributed to the polyethoxylated castor oil (Cremophor EL) vehicle (Tammisto et al. 1973) made its use redundant. Subsequent formulations of alfaxalone incorporating a cyclodextrin have hitherto been devoid of the previous side effects and Alfaxan (alfaxalone dissolved in 2-hydroxypropyl- β -cyclodextrin) is now registered for induction and maintenance of anaesthesia in dogs and cats and has been used in horses (Goodwin et al. 2011) sheep (Andaluz et al. 2012; del Mar Granados et al. 2012) rabbits (Navarrete-Calvo et al. 2014) and other more exotic species (Bouts & Karunaratna 2011; McMillan & Leece 2011; Bauquier et al. 2013; Kischinovsky et al. 2013; Knotek et al. 2013; Villaverde-Morcillo et al. 2014). The use of alfaxalone in biomedical research and clinical veterinary medicine is gaining popularity as it may offer some selective advantages over other anaesthetic combinations in terms of safety, reflex suppression, cardiopulmonary depression, interaction with receptors involved in pain pathways/modulation and pain on injection (Child et al. 1972; Michou et al. 2012; Santos González et al. 2013) but may also offer additional advantages in influencing CNS development and myelination (Yawno et al.

2014). Alfaxalone has been popular for neuroendocrine studies for its sparing of various forebrain functions (Sarkar et al. 1976; Sherwood et al. 1980). Human trials of alphaxalone in cyclodextrin are currently underway (Monagle et al. 2015). The majority of animals used in basic science pain research however are young healthy male laboratory rodents, and indeed it has been suggested that a more heterogeneous and diverse population be used to improve the translational relevance to a human population (Mogil 2009). The inclusion of female rodents is to be encouraged despite the additional complexities that the variability of the oestrous cycle and sexual dimorphism poses; well-designed studies can include both sexes without needless increase in animal numbers (Clayton & Collins 2014). With respect to alfaxalone usage, the pharmacokinetics of a single intravenous (IV) dose have been defined in dogs (Ferré et al. 2006) cats (Whittem et al. 2008b; Muir et al. 2009) female rats (Lau et al. 2013) and male rats after a 5 minute infusion (Visser et al. 2002). The novelty and primary aim of this study was therefore to characterise the pharmacokinetics, pharmacodynamics and hypnotic characteristics of a constant rate infusion of alfaxalone in male *versus* female rats.

Materials and methods

This study was performed in accordance with Project Licence PPL30/3156 issued under the Animal (Scientific) Procedures Act 2013 (EU Directive 2010/63/EU) and local ethics committee as part of a larger study investigating nociceptive withdrawal reflexes and diffuse noxious inhibitory control. This study is reported in accordance with the ARRIVE guidelines (Kilkenny et al. 2014).

91 Animals

92 Twelve (9-12 week old) Sprague Dawley rats, six male (397 ± 16 g) and six female
93 (286 ± 20 g) (Charles River Laboratories, Margate, UK) were used. Animals were
94 housed in single sex groups of four, given access to food (Teklad 2018, Harlan) and tap
95 water *ad libitum* and maintained on a 12-hour light/dark cycle. All experiments started
96 at 10:00 h each day.

97

98 General anaesthesia

99 Anaesthesia was induced using 3% isoflurane (Isoflo; Abbott, UK) in oxygen and
100 nitrous oxide mixture (1:2). Once the rat had lost its righting reflex, it was transferred to
101 a heating blanket (Harvard Apparatus Ltd., UK) coupled to a rectal probe for
102 maintenance of body temperature ($37.5 \pm 0.5^\circ\text{C}$). Anaesthesia was maintained using
103 2.00-2.25% (vaporizer setting) isoflurane in oxygen/nitrous oxide delivered via a
104 nosecone. Lidocaine 2% (Lignol; Dechra, UK) 3 mg kg^{-1} was infiltrated subcutaneously
105 prior to skin and sternohyoid incision. The trachea was surgically cannulated using
106 polyethylene 2.42mm O.D. tubing (Fisher Scientific, UK). Respiratory rate and effort
107 was assessed by observing chest excursion and measuring end tidal carbon dioxide
108 (CapStar 100, Linton, Diss, UK). In animals exhibiting respiratory depression as judged
109 by a low respiratory rate and rising end tidal carbon dioxide values, intermittent positive
110 pressure ventilation was initiated (Harvard 683 ventilator, Harvard Apparatus, UK) at
111 60-80 breaths minute^{-1} to maintain end tidal carbon dioxide at 35-45 mmHg (4.67-6.00
112 kPa). The left jugular vein was surgically cannulated using 0.63 mm O.D. polyethylene
113 tubing (Fisher Scientific) for administration of alfaxalone. The left carotid artery was

114 surgically cannulated using 1mm O.D. polyethylene tubing (Fisher Scientific) to
115 monitor arterial blood pressure and for sampling. Arterial blood pressure was monitored
116 by an arterial pressure transducer (Sensonor 840; SensoNor, Norway) and recorded
117 using a PC running Spike2 software (CED Ltd, UK). Heart rate was recorded via two 25
118 gauge needles inserted subcutaneously on the lateral sides of the thoracic wall. The
119 electrocardiogram signal was amplified and used to trigger an instant rate meter
120 (Neurolog NL253, Digitimer, UK) and again recorded using Spike2 software.
121 An infusion of alfaxalone (Alfaxan, Jurox, UK) was started at time 0 at a loading dose
122 of $1.67 \text{ mg kg}^{-1} \text{ minute}^{-1}$ for 2.5 minutes followed by a constant rate infusion (0.75 mg
123 $\text{kg}^{-1} \text{ minute}^{-1}$) for the remainder of the electrophysiological experiment using a
124 calibrated syringe driver (SP100iz, WPI, UK). The isoflurane and nitrous oxide were
125 stopped 2.5 minutes after starting the alfaxalone infusion. Arterial blood was withdrawn
126 from the carotid cannula into lithium heparin and placed on ice. Blood samples (200 μl)
127 were collected at baseline (prior to alfaxalone), end of loading dose, and at 10, 30, 60,
128 90, 120 minutes. Arterial blood gases, biochemistry and haematology parameters (pH,
129 pCO_2 , pO_2 , bicarbonate, sodium, potassium, chloride, calcium, glucose, lactate and
130 creatinine concentrations) were also measured (EPOC, Woodley Instrumentation,
131 Bolton, Lancashire, UK). All rats received an equal volume of balanced electrolyte
132 solution after sampling (Vetivex 11 (Hartmann's); Dechra, UK). Samples were
133 centrifuged (4000g for 10 minutes) within 30 minutes of collection. Plasma was
134 harvested and stored at -20°C until determination of plasma alfaxalone concentration.
135 The hypnotic characteristics of the anaesthetic were evaluated by monitoring paw
136 withdrawal reflex in response to pinch, corneal reflex in response to light brushing,

spontaneous blinking and gross purposeful movement and cardiopulmonary parameters. Following the pharmacokinetic study, a separate electrophysiological study was performed, EMG responses were recorded from tibialis anterior, biceps femoris, and medial gastrocnemius muscles during electrical plantar hind paw stimulation of the toes and heel, before and after a conditioning injection of capsaicin into either the contralateral forelimb to study diffuse noxious inhibitory controls (DNIC) or the ipsilateral hind limb to investigate central sensitization and reflex facilitation (Harris & Clarke 2003). Data from this part of the study were not included in this paper but informed the subjective assessment of response to noxious stimuli during alfaxalone anaesthesia. At the end of the experiments animals were euthanised by IV injection of pentobarbitone (Pentobarbital; Ayrton Saunders Ltd, UK) followed by cervical dislocation (as required by UK Home Office regulations). All female rats underwent vaginal swabbing to characterise vaginal smear cell types. Slides were examined under x40 and then x100 magnifications (BH2 microscope, Olympus, UK) after staining with modified Giemsa (Diff Quik, Vet Direct, UK) and cell types and numbers were recorded.

Sample analyses

Samples were analyzed for alfaxalone using a LCMS/MS method. Methanolic standard curve and quality control (QC) spiking solutions were produced for alfaxalone from separate accurate weighings of solid compound. Standards and QCs were prepared by spiking 10 µl spike solution into a solution of 20 µl plasma + 30 µl water + 40 µl methanol + 150 µl methanol containing 1000 nM tolbutamide as internal standard.

Plasma standard curves were prepared from 100 – 5000 ng mL⁻¹ and QCs were prepared for 250 and 2500 ng mL⁻¹. Blank male or female plasma was used for the standards and QC solutions (Charles River, UK). The plasma samples were prepared by adding 30 µl water + 50 µl methanol + 150 µl methanol containing 1000 nM tolbutamide as internal standard to 20 µl plasma. Samples, standards and QCs were then mixed and stored in a freezer at -20°C for a minimum of 120 minutes prior to centrifugation at 4000g for 20 minutes. The samples were extracted and analyzed using a Micromass Quattro Premier mass spectrometer incorporating an Acquity autosampler (Waters, UK). An ACE Excel 2 C18-AR 50 x 2.1mm column was used with the following LC conditions: Solvent A = Water + 0.1% Formic Acid, Solvent B = Methanol + 0.1% Formic Acid, Flow rate = 0.8 mL minute⁻¹, column temperature = 60°C. LC gradient went from 95 % solvent A:5 % solvent B to 5 % solvent A:95 % solvent B over a 1.5 minute interval. The MS/MS method used electrospray positive mode with a 333.16 >107.01 transition for the detection of alfaxalone. The lower limit of quantification (LLOQ) was 100 ng mL⁻¹. The coefficient of variation at LLOQ was <8% and <16% for other concentration levels. All samples were run in triplicate. Two separate LC/MS/MS runs were performed for the male and female samples, respectively.

Samples were analysed within 28 days of collection based on data from analytical validation study file supporting stability of alfaxalone in rat plasma at -20 ° for 30 days (Jurox 2010).

Pharmacokinetic analyses

Pharmacokinetic analysis was carried out using Phoenix WinNonlin 6.3 (Pharsight, Sunnyvale, CA, USA). The pharmacokinetic parameters (clearance, volume of distribution and half life) for each individual rat were estimated according to best fit from an IV infusion one compartmental pharmacokinetic model, based on previous published data (Lau et al. 2013) showing a single exponential decay, with appropriate weighting for best fit. The appropriate weighting for best fit in this study was based on examination of the residuals showing random scatter around predicted values using $1/y^2$ weighting by 1/reciprocal of the predicated value.

Statistical Analyses

Statistical tests were performed using GraphPrism (GraphPad Software, CA, USA) version 6. The pharmacokinetic parameters for log transformed parameter data for both genders were compared using an unpaired, two tailed Student's *t*-test and a *p* value of <0.05 was considered significant. Data are reported as mean \pm standard deviation (SD) unless stated otherwise.

Cardiopulmonary data were collected continuously, and analyzed at the pharmacokinetic time points. The normality assumptions were tested with Kolmogorov-Smirnov or Shapiro-Wilk tests. The differences in heart rate, mean/systolic/diastolic arterial blood pressure, blood gas variables, lactate, glucose and electrolytes between genders were compared using an unpaired *t*-test. For a more detailed analysis of changes over time, these variables were also analyzed by two-way repeated-measures ANOVA (one factor repetition) for the time points between baseline anaesthesia

and 120 minutes after alfaxalone infusion with Sidak's correction for multiple comparisons.

Results

Anaesthetic induction and instrumentation were completed without difficulty in all animals. All animals underwent a total of 230 ± 20 minutes of alfaxalone anaesthesia consisting of a 120 minutes of a PK/PD study followed by an electrophysiology study. Only results from the former are reported here.

Pharmacokinetics

The shape of the concentration-time curve following a loading dose and then constant rate infusion was typical of those observed for anaesthetic induction drugs exhibiting an initial steep phase after the loading dose followed by a gradual increase until steady state was achieved. The plasma concentrations were substantially different between the sexes (Figure 1).

The pharmacokinetic parameters calculated by an IV infusion one compartmental model are shown in Table 1. Logarithmic transformed data for clearance and $t_{1/2}$ was significantly different between the male and female rats by the two-tailed t -test. As would be expected the MRT was also significantly different between genders.

Pharmacodynamics

Cardiopulmonary data are presented in Table 2.

226 After a 2.5 minute loading dose all rats showed an initial short lived decrease in arterial
227 blood pressure, heart rate and respiratory rate as a result of concomitant administration
228 of inhalant and alfaxalone. Within the next 5 minutes following discontinuation of
229 isoflurane, all rats demonstrated an increase in blood pressure from baseline reading
230 under isoflurane and nitrous oxide anaesthesia. Blood pressure (mean, systolic,
231 diastolic), heart rate and respiratory rate at baseline were not significantly different
232 between male and female rats under isoflurane anaesthesia.

233 Heart rates remained stable during alfaxalone anaesthesia and there was no significant
234 difference between the sexes at any time points.

235 Systolic, mean and diastolic arterial pressures all increased from baseline under
236 isoflurane anaesthesia, reached a peak (between 60 and 90 minutes) and thereafter
237 showed a trend of decreasing with time. Mean arterial blood pressure was significantly
238 different between males and female rats ($p = 0.026$), however the interaction with time
239 ($p < 0.0001$) differed between the genders for mean (and systolic and diastolic) pressure
240 and makes the interpretation of these data difficult (Fig. 2). Significant differences were
241 analyzed *post hoc* using Sidak's multiple comparison test. Mean arterial blood pressure
242 was significantly increased compared to baseline in male rats at 30, 60, 90 and 120
243 minutes from starting the alfaxalone ($p < 0.0005$), as were systolic and diastolic
244 pressures. Mean arterial blood pressure was only significantly greater than baseline in
245 the female rats at 60 minutes. In 4 of 12 animals (2 males and 2 females)
246 cardiopulmonary depression, indicated by a decrease in blood pressure or respiratory
247 rate, necessitated discontinuation of the isoflurane and nitrous oxide before the 2.5

minute time point. Within 60 seconds of discontinuation of the isoflurane and nitrous oxide, heart rate and blood pressure began to rise in all animals.

All female rats ventilated spontaneously throughout the experiment, whereas 2 of 6 male rats required mechanical ventilation as judged by apnoea, or a rise in end tidal carbon dioxide coupled with a decrease in respiratory rate and effort.

Blood gas parameters and biochemistry values are presented in Table 3. There were no significant differences between sexes for these parameters except for pH ($p = 0.0027$), which was lower in the female rats in conjunction with higher partial pressures of carbon dioxide. The clinical significance of this is unknown and of little significance.

Partial pressures of oxygen were different between baseline and subsequent time points, trending towards higher values under total intravenous anaesthesia compared to inhalational anaesthesia.

Hypnotic effect

The plane of anaesthesia was continually evaluated by serial cardiopulmonary measurements, blood gas analysis and reflex responses. Subjective evaluation of this hypnotic effect of the alfaxalone in all 12 rats was excellent. No rats demonstrated gross purposeful movement or required a change in the infusion rate to improve the plane of anaesthesia.

Vaginal smears

The same investigator read all slides (3 per rat) and evaluated the whole slide to give an impression of the smear, rather than exact cell counts. Three rats were characterized as in dioestrus, one in proestrus, one in oestrus, and one in metoestrus.

Discussion

The major finding from this study is that consideration must be given to the dose of anaesthetic delivered to male and female subjects. Without this interrogation of the experimental model there remains the danger that studies will be carried out under what is assumed to be identical 'planes' of anaesthesia, when in reality one sex may be more or less profoundly anaesthetized such that, for example, hormonal or neuroendocrine responses will be affected.

The data reported here were part of an electrophysiological study investigating diffuse noxious inhibitory controls of nociceptive withdrawal reflexes and is part of a larger study of descending control in chronic osteoarthritis, which in humans is more prevalent in females, and highlights the potential risk that using one sex may contribute to the failure of Phase 1 trials or misleading conclusions. In many studies, the influence of the anaesthetic is ignored, or so poorly reported that ensuring a consistent plane of anaesthesia is impossible. Although rats are frequently used in laboratory studies involving anaesthesia, it is typically males used to reduce experimental variability (Zucker & Beery 2010). The limitation of this approach is that basic science intended to be translated into the human population is potentially compromised by the use of one sex (Clayton & Collins 2014). This perceived variability in females is often used as a reason for excluding females from studies. A meta-analysis of 293 studies in which murine behavioural, morphological, physiological, and molecular traits were monitored

in both sexes showed variability was not significantly greater in females for any endpoint but several traits contributed to substantially greater variability in the males in this analysis, including the influence of group housing (Prendergast et al. 2014).

Commendably, there is a movement towards trying to include more female subjects in studies.

The small sample size in our study makes it impossible to postulate that differences within the females are a result of the differences in the oestrous cycle or as a result of normal variability; nonetheless it demonstrates that if steroid hormones can affect alfaxalone efficacy, the stage of the oestrous cycle may also contribute. It is possible that the three female rats requiring additional isoflurane for several minutes between minute 6 and 10 after commencing the alfaxalone may be a result of the stage of the oestrous cycle, but this is impossible to prove definitively.

In the research community there remains a collective responsibility for a thoroughness of reporting anaesthesia conditions in order that the anaesthesia is not ‘the elephant in the room’.^a

Other species e.g. cats display similar pharmacokinetic characteristics to this rat study (Whittem et al. 2008a). The characteristic rapid hepatic metabolic clearance of alfaxalone by the liver has been identified *in-vivo* and *in-vitro* in rats (Sear & McGivan

^a “**Elephant in the room**” is an English metaphorical idiom for an obvious truth that is either being ignored or going unaddressed. The idiomatic expression also applies to an obvious problem or risk no one wants to discuss.

1980; Sear & McGivan 1981). Gender based differences in drug metabolism are the primary cause of sex-dependent pharmacokinetics and reflect differences in the expression of hepatic enzymes active in the metabolism of many extrinsic and intrinsic chemicals, including cytochrome P450 (Waxman & Holloway 2009). Rodent studies have identified more than a 1000 genes whose expression is dependent on sex and these genes modulate liver metabolic function and create sexual dimorphism in liver function (Tanaka 1999). Differences in bioavailability, distribution, metabolism, and/or excretion in different sexes are multifactorial and complicated (Soldin & Mattison 2009). Drug distribution can also be sex linked, influenced by factors such as body fat, plasma volume and differential perfusion of organs. However, in this study no significant difference was recorded between male and female V_{ss} suggesting this was not an issue. In general, however, sex differences in metabolism are thought to be the primary determinant of variation in pharmacokinetics and this is most likely the reason for the differences seen in this study.

The quality of anaesthesia was subjectively judged as excellent in all rats. Contrary to previous studies (Brammer et al. 1993), this anaesthetic combination provided very good conditions, stability and survival beyond 180 minutes. The lower blood pressures observed in female rats in this study would suggest that these animals were more profoundly anaesthetized than the males. There was no difference in heart rates between the groups, and the depth of anaesthesia was not so sufficiently profound as to cause apnoea in the females. This is likely to be a pharmacokinetic effect as the females' clearance of alfaxalone is so much less than in the male rats. The consistency of the

haematology, biochemistry and blood gas values in all rats demonstrates the stability of the protocol. Blood gas values are infrequently reported for rodent anaesthetics, in part due to the technical nature of artery cannulation, and previously the volume of blood required made repeated sampling impossible due to a deleterious depletion of the blood volume of the animal. Newer point of care analyzers are able to process much smaller volumes. Total blood volume removed was well below the limit of 10% blood volume and this removal of blood in conjunction with a replenishment of balanced electrolyte solution clearly had no impact on the animals. All biochemistry and haematology values except chloride were similar to those provided by the supplier of the Sprague Dawley rats in age-matched subjects. Invariably these samples were analysed with different machines, but even with slight discrepancies usually seen between laboratories, plasma chloride values in the study (mmol L^{-1}) (114 ± 2.6 (females) 115 ± 2.7 (males)) and values in age matched conscious Sprague Dawley rats (109 ± 1.4 (females), 103 ± 1.00 (males) were different. A moderate corrected hyperchloraemia was present (when measured sodium values were also taken into consideration). The most likely cause was the administration of normal heparinized saline during cannula placement and through flushing of the carotid cannula with heparinized saline to maintain patency for sampling and blood pressure measurement. A concurrent acidosis was not observed, and sodium values were almost identical between the supplied conscious values and those measured. The potential deleterious effects of normal saline administration have been raised (Handy & Soni 2008) and the administration of non-physiological saline and balanced electrolyte solutions is warranted. The clinical impact of hyperchloraemia is unknown in this study, but in humans there is an increasing awareness that

hyperchloraemia and hyperchloraemic acidosis can cause significant clinical ramifications (Handy & Soni 2008).

Differences in efficacy of the older alfaxalone steroid (Alphathesin/Althesin) in male and female rats has been demonstrated, with males requiring four times the dose of females for surgical anaesthesia and analgesia, and it was concluded that the influence of sex hormones was responsible for this discrepancy (Fink et al. 1982). However a recent study disputed this and postulated that differences are more likely a result of the different formulations of alfaxalone and assay methodologies than differences between sexes (Lau et al. 2013). Notwithstanding these views it has also been shown that formulation is hugely influential; the toxicity of alfaxalone in Wistar rats was less in those animals receiving alfaxalone dissolved in 7-sulfobutyl-ether- β -cyclodextrin (SBECD) compared to alfaxalone in Cremophor EL (Goodchild et al. 2015). The current study directly comparing male and female Sprague Dawley rats receiving an HPCD alfaxan formulation seemingly favours a true sex difference due to pharmacokinetics, pharmacodynamics or both as the explanation. Sex based studies using rats anaesthetized with pentobarbitone have also demonstrated differences (Zambricki & Dalecy 2004). Sex differences have been detected in studies comparing IP and IV routes in rats (Estes et al. 1990) postulating that the lack of obvious sex differences with single IV dosing may be a result of the short duration of effect. Group sizes of 6 were deemed appropriate for evaluating a drug exhibiting within subject variability of less than 30% coefficient of variation of pharmacokinetic measures (Rowland & Tozer 2011) and recent studies comparing cardiovascular effects

of anaesthetic drugs have successfully used 5 rats per group (Bencze et al. 2013). It should also be noted that the presence of isoflurane and nitrous oxide at the outset of the loading dose is likely to have affected cardiopulmonary parameters in the very early stages of the infusion and conclusions drawn about differences at these time points are likely to be tenuous. However both groups underwent identical protocols so this influence would have been similar for both groups. The use of an inhalational agent such as isoflurane with minimal metabolism and rapid elimination ensured that the period of time from ceasing administration was as short as possible. The maximum possible duration of initial concurrent administration was 2.5 minutes.

In summary, there are pharmacokinetic and pharmacodynamic differences with alfaxalone in cyclodextrin in male and female rats. The plane of anaesthesia provided by this protocol is stable and clinically indistinguishable between sexes with no apparent cumulative effect. Half-life, clearance and mean residence time were significantly different between male and female rats indicating that a sex-linked effect was present. The protocol in our study provides excellent anaesthesia conditions but concludes that a dose alteration may be necessary for rat sex-based studies incorporating alfaxalone. This contrasts to previous published studies, which have dismissed a sex difference (Ferré et al. 2006; Berry 2015). Population pharmacokinetics are necessary to further investigate these findings.

Acknowledgements

Victoria Simmonds for technical assistance.

403 This study received alfaxalone (Alfaxan) donated by Jurox, Malvern, UK. No conflicts
404 of interest are declared.

405 **Authors' contributions**

406 KLW, JH: study design and planning; KLW, JH: study conduct; KLW, SWP: data
407 analysis; all authors: paper writing.

408

References

- Andaluz A, Felez-Ocaña N, Santos L et al. (2012) The effects on cardio-respiratory and acid-base variables of the anaesthetic alfaxalone in a 2-hydroxypropyl- β -cyclodextrin (HPCD) formulation in sheep. *The Veterinary journal* 191, 389-392.
- Bauquier SH, Greenwood J, Whittem T (2013) Evaluation of the sedative and anaesthetic effects of five different concentrations of alfaxalone in goldfish, *Carassius auratus*. *Aquaculture* 396-399, 119-123.
- Bencze M, Behuliak M, Zicha J (2013) The impact of four different classes of anesthetics on the mechanisms of blood pressure regulation in normotensive and spontaneously hypertensive rats. *Physiol Res* 62, 471-478.
- Berry SH (2015) *Injectable Anesthetics*. (5th edn). Grimm KA, Lamont LA, Tranquilli WJ, et al. (eds). Wiley Blackwell, Ames, Iowa. pp. 277-296.
- Bouts T, Karunaratna D (2011) Evaluation of medetomidine-alfaxalone and medetomidine-ketamine in semi-free ranging Bennett's wallabies (*Macropus rufogriseus*). *Journal of Zoo and Wildlife Medicine* 42, 617-622.
- Brammer a, West CD, Allen SL (1993) A comparison of propofol with other injectable anaesthetics in a rat model for measuring cardiovascular parameters. *Laboratory animals* 27, 250-257.
- Child KJ, Davis B, Dodds MG et al. (1972) Anaesthetic, cardiovascular and respiratory effects of a new steroidal agent CT 1341: a comparison with other intravenous anaesthetic drugs in the unrestrained cat. *British journal of pharmacology* 46, 189-200.

- 432 Clayton JA, Collins FS (2014) Policy: NIH to balance sex in cell and animal studies.
433 Nature 509, 282-283.
- 434 del Mar Granados M, Dominguez JM, Fernandez-Sarmiento A et al. (2012) Anaesthetic
435 and cardiorespiratory effects of a constant-rate infusion of alfaxalone in
436 desflurane-anaesthetised sheep. pp. 125-125.
- 437 Estes KS, Brewster ME, Webb AI et al. (1990) A non- surfactant formulation for
438 alfaxalone based on an amorphous cyclodextrin: Activity studies in rats and
439 dogs. International Journal of Pharmaceutics 65, 101-107.
- 440 Ferré PJ, Pasloske K, Whitem T et al. (2006) Plasma pharmacokinetics of alfaxalone in
441 dogs after an intravenous bolus of Alfaxan-CD RTU. Veterinary anaesthesia and
442 analgesia 33, 229-236.
- 443 Fink G, Sarkar DK, Dow RC et al. (1982) Sex difference in response to alphaxalone
444 anaesthesia may be oestrogen dependent. Nature 298, 270-272.
- 445 Goodchild CS, Serrao JM, Kolosov A et al. (2015) Alphaxalone Reformulated: A
446 Water-Soluble Intravenous Anesthetic Preparation in Sulfobutyl-Ether-beta-
447 Cyclodextrin. Anesth Analg 120, 1025-1031.
- 448 Goodwin Wa, Keates HL, Pasloske K et al. (2011) The pharmacokinetics and
449 pharmacodynamics of the injectable anaesthetic alfaxalone in the horse.
450 Veterinary anaesthesia and analgesia 38, 431-438.
- 451 Handy JM, Soni N (2008) Physiological effects of hyperchloraemia and acidosis.
452 British journal of anaesthesia 101, 141-150.
- 453 Harris J, Clarke RW (2003) Organisation of sensitisation of hind limb withdrawal
454 reflexes from acute noxious stimuli in the rabbit. J Physiol 546, 251-265.

- 455 Harrison N, Simmonds M (1984) Modulation of the GABA receptor complex by a
456 steroid anesthetic. *Brain research* 323, 287-292.
- 457 Jurox (2010) Study Report JX9604.03-V027- Validation of an LC/MS Analytical
458 Method for the Analysis of Alfaxalone in Rat Plasma.
- 459 Kilkenny C, Browne W, Cuthill I et al. (2014) Improving Bioscience Research
460 Reporting: The ARRIVE Guidelines for Reporting Animal Research. *Animals* 4,
461 35-44.
- 462 Kischinovsky M, Duse A, Wang T et al. (2013) Intramuscular administration of
463 alfaxalone in red-eared sliders (*Trachemys scripta elegans*) - effects of dose and
464 body temperature. *Veterinary anaesthesia and analgesia* 40, 13-20.
- 465 Knotek Z, Hrdá A, Knotková Z et al. (2013) Alfaxalone anaesthesia in the green iguana
466 (*Iguana iguana*). *Acta Veterinaria Brno* 82, 109-114.
- 467 Lau C, Ranasinghe MG, Shiels I et al. (2013) Plasma pharmacokinetics of alfaxalone
468 after a single intraperitoneal or intravenous injection of Alfaxan in rats. *J Vet*
469 *Pharmacol Ther* 36, 516-520.
- 470 McMillan MW, Leece EA (2011) Immersion and branchial/transcutaneous irrigation
471 anaesthesia with alfaxalone in a Mexican axolotl. *Veterinary anaesthesia and*
472 *analgesia* 38, 619-623.
- 473 Michou JN, Leece EA, Brearley JC (2012) Comparison of pain on injection during
474 induction of anaesthesia with alfaxalone and two formulations of propofol in
475 dogs. *Veterinary anaesthesia and analgesia* 39, 275-281.
- 476 Mogil JS (2009) Animal models of pain: progress and challenges. *Nat Rev Neurosci* 10,
477 283-294.

- 478 Monagle J, Siu L, Worrell J et al. (2015) A Phase 1c Trial Comparing the Efficacy and
479 Safety of a New Aqueous Formulation of Alphaxalone with Propofol. *Anesth*
480 *Analg* 121, 914-924.
- 481 Muir W, Lerche P, Wiese A et al. (2009) The cardiorespiratory and anesthetic effects of
482 clinical and supraclinical doses of alfaxalone in cats. *Veterinary anaesthesia and*
483 *analgesia* 36, 42-54.
- 484 Navarrete-Calvo R, Gómez-Villamandos RJ, Morgaz J et al. (2014) Cardiorespiratory,
485 anaesthetic and recovery effects of morphine combined with medetomidine and
486 alfaxalone in rabbits. *The Veterinary record* 174, 95-95.
- 487 Prendergast BJ, Onishi KG, Zucker I (2014) Female mice liberated for inclusion in
488 neuroscience and biomedical research. *Neuroscience and Biobehavioral*
489 *Reviews* 40, 1-5.
- 490 Rowland M, Tozer T (2011) Variability. In: *Clinical Pharmacokinetics and*
491 *Pharmacodynamics: Concepts and Applications*. (4th Ed edn). Lippincott
492 Williams & Wilkins, Philadelphia. pp. 333-355.
- 493 Rupprecht R, Holsboer F (1999) Neuroactive steroids: mechanisms of action and
494 neuropsychopharmacological perspectives. *Trends in neurosciences* 22, 410-
495 416.
- 496 Santos González M, Bertrán de Lis BT, Tendillo Cortijo FJ (2013) Effects of
497 intramuscular alfaxalone alone or in combination with diazepam in swine.
498 *Veterinary anaesthesia and analgesia* 40, 399-402.
- 499 Sarkar DK, Chiappa SA, Fink G et al. (1976) Gonadotropin-releasing hormone surge in
500 pro-oestrous rats. *Nature* 264, 469-463.

- 501 Sear JW, McGivan JD (1980) The metabolism of alphaxalone by isolated rat
502 hepatocytes. *Biochem Pharmacol* 29, 248-248.
- 503 Sear JW, McGivan JD (1981) Metabolism of Alphaxalone in the Rat: Evidence for the
504 Limitation of the Anaesthetic Effect By the Rate of Degradation Through the
505 Hepatic Mixed Function Oxygenase System. *British journal of anaesthesia* 53,
506 417-424.
- 507 Sherwood NM, Chiappa SA, Sarkar DK et al. (1980) Gonadotropin-Releasing Hormone
508 (GnRH) in Pituitary Stalk Blood from Proestrous Rats: Effects of Anesthetics
509 and Relationship Between Stored and Released GnRH and Luteinizing
510 Hormone. *Endocrinology* 107, 1410-1417.
- 511 Soldin OP, Mattison DR (2009) Sex Differences in Pharmacokinetics and
512 Pharmacodynamics. *Clinical Pharmacokinetics* 48, 143-157.
- 513 Tammisto T, Takki S, Tigerstedt I et al. (1973) A comparison of althesin and
514 thiopentone in induction of anaesthesia. *British journal of anaesthesia* 45, 100-
515 107.
- 516 Tanaka E (1999) Gender-related differences in pharmacokinetics and their clinical
517 significance. *Journal of clinical pharmacy and therapeutics* 24, 339-346.
- 518 Turner DM, Ransom RW, Yang JS et al. (1989) Steroid anesthetics and naturally
519 occurring analogs modulate the gamma-aminobutyric acid receptor complex at a
520 site distinct from barbiturates. *The Journal of pharmacology and experimental*
521 *therapeutics* 248, 960-966.
- 522 Villaverde-Morcillo S, Benito J, García-Sánchez R et al. (2014) Comparison of
523 isoflurane and alfaxalone (Alfaxan) for the induction of anesthesia in flamingos

- 524 (Phoenicopterus roseus) undergoing orthopedic surgery. Journal of zoo and
525 wildlife medicine 45, 361-366.
- 526 Visser SAG, Smulders CJGM, Reijers BPR et al. (2002) Mechanism-based
527 pharmacokinetic-pharmacodynamic modeling of concentration-dependent
528 hysteresis and biphasic electroencephalogram effects of alphaxalone in rats. The
529 Journal of pharmacology and experimental therapeutics 302, 1158-1167.
- 530 Waxman DJ, Holloway MG (2009) Sex Differences in the Expression of Hepatic Drug
531 Metabolizing Enzymes. Mol Pharmacol 76, 215-228.
- 532 Whittem T, Pasloske KS, Heit MC et al. (2008a) The pharmacokinetics and
533 pharmacodynamics of alfaxalone in cats after single and multiple intravenous
534 administration of Alfaxan at clinical and supraclinical doses. Journal of
535 veterinary pharmacology and therapeutics 31, 571-579.
- 536 Whittem T, Pasloske KS, Heit MC et al. (2008b) The pharmacokinetics and
537 pharmacodynamics of alfaxalone in cats after single and multiple intravenous
538 administration of Alfaxan at clinical and supraclinical doses. Journal of
539 Veterinary Pharmacology and Therapeutics 31, 571-579.
- 540 Yawno T, Mortale M, Sutherland AE et al. (2014) The effects of betamethasone on
541 allopregnanolone concentrations and brain development in preterm fetal sheep.
542 Neuropharmacology 85, 342-348.
- 543 Zambricki EA, Dalecy LG (2004) Rat sex differences in anesthesia. Comp Med 54, 49-
544 53.
- 545 Zucker I, Beery AK (2010) Males still dominate animal studies. Nature 465, 690-690.
546

547

548

Figure Legends

Figure 1 Mean (\pm SD) alfaxalone plasma concentrations (ng mL^{-1}) for 12 Sprague Dawley rats (6 female, 6 male) after intravenous administration of alfaxalone $1.67 \text{ mg kg}^{-1} \text{ minute}^{-1}$ for 2.5 minutes followed by $0.75 \text{ mg kg}^{-1} \text{ minute}^{-1}$ alfaxalone for 120 minutes using a one compartment infusion model.

Figure 2 Mean (\pm SD) arterial blood pressure (MAP) (mmHg) for 12 Sprague Dawley rats (6 female, 6 male) after intravenous administration of alfaxalone $1.67 \text{ mg kg}^{-1} \text{ minute}^{-1}$ for 2.5 minutes followed by $0.75 \text{ mg kg}^{-1} \text{ minute}^{-1}$ alfaxalone for 120 minutes using a one compartment infusion model.

Table 1 Pharmacokinetic parameters for 12 Sprague Dawley rats (6 female, 6 male) after intravenous administration of alfaxalone at a rate of $1.67 \text{ mg kg}^{-1} \text{ minute}^{-1}$ for 2.5 minutes followed by $0.75 \text{ mg kg}^{-1} \text{ minute}^{-1}$ for 120 minutes using a one compartment infusion model.

Rat ID	CL (mL minute ⁻¹ kg ⁻¹)	T _{1/2}	Vdss L kg ⁻¹	MRT minutes	Cmax mg L ⁻¹
Female 1	66.3	33.8	3.24	48.8	1.3
Female 2	40.4	29.8	1.74	43.0	2.3
Female 3	46.9	76.7	5.19	110.7	0.8
Female 4	58.2	30.9	2.60	44.6	1.6
Female 5	31.9	69.2	3.19	99.9	1.3
Female 6	42.3	34.1	2.08	49.2	2.0
Mean	47.7	45.8	3.00	66.0	1.5
Sd	8.86	20.7	0.6	30.7	0.6
Male 1	65.8	52.1	4.94	75.1	0.8
Male 2	79.9	13.2	1.53	19.1	2.6
Male 3	78.5	17.3	1.95	24.9	2.0
Male 4	117.9	10.3	1.75	14.8	2.2
Male 5	106.2	18.6	2.85	26.8	1.4

Male 6	101.2	26.5	3.87	38.3	1.0
Mean	91.6	23.0	2.82	33.2	1.7
Sd	19.9	15.3	1.36	22.0	0.7
P value	0.0008***	0.0268*	0.710	0.027*	0.780

568

569

570 CL = clearance, $t_{1/2}$ = half life, V_{dss} = volume of distribution, MRT = mean residence
 571 time, C_{max} = maximum plasma concentration

572 CL, $t_{1/2}$ and MRT is significantly different between the male and female rats.

573 Asterisks denote significant difference * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

574

Table 2 Cardiopulmonary parameters for 12 Sprague Dawley rats (6 female, 6 male) after intravenous administration of alfaxalone at a rate of $1.67 \text{ mg kg}^{-1} \text{ minute}^{-1}$ for 2.5 minutes followed by $0.75 \text{ mg kg}^{-1} \text{ minute}^{-1}$ for 120 minutes using a one compartment infusion model.

Variable	Sex	-5	2.5	10	30	60	90	120
MAP	Male	97 ± 7	71 ± 7	$110 \pm 32^*$	$127 \pm 17^{*****}$	$138 \pm 11^{*****}$	$135 \pm 16^{*****}$	$132 \pm 9^{*****}$
	Female	89 ± 13	90 ± 20	110 ± 17	114 ± 18	$121 \pm 14^{**}$	113 ± 23	114 ± 21
SAP	Male	98 ± 8	96 ± 9	$132 \pm 33^{**}$	$153 \pm 22^{*****}$	$167 \pm 17^{*****}$	$167 \pm 17^{*****}$	$158 \pm 11^{*****}$
	Female	108 ± 12	111 ± 17	127 ± 14	133 ± 15	$144 \pm 15^{**}$	$138 \pm 25^*$	$138 \pm 24^*$
DAP	Male	69 ± 7	60 ± 9	$98 \pm 32^*$	$114 \pm 15^{***}$	$124 \pm 11^{*****}$	$119 \pm 19^{*****}$	$120 \pm 8^{*****}$
	Female	79 ± 14	79 ± 21	101 ± 19	104 ± 19	$109 \pm 14^*$	100 ± 22	101 ± 21
HR	Male	$440 \pm$	$442 \pm$	440 ± 15	455 ± 23	447 ± 28	438 ± 34	423 ± 23

		25	28					
RR	Female	440 ±	440 ±	436 ± 35	455 ± 23	445 ± 22	415 ± 12	422 ± 16
		31	31					
	Male	58 ± 12	55 ± 10	56 ± 10	68 ± 10	57 ± 14	65 ± 7	71 ± 9
	Female	56 ± 10	54 ± 7	56 ± 6	55 ± 5	63 ± 5	68 ± 4	69 ± 4

MAP: mean arterial pressure (mmHg); SAP: systolic arterial pressure (mmHg); DAP: diastolic arterial pressure (mmHg); HR: heart rate (beats per minute); RR: respiratory rate (breaths per minute)

Baseline (-5 minute) samples were all measured under isoflurane (2% vaporizer setting) in N₂O and O₂. Data are mean ± SD.

Asterisks denote significant difference from baseline (-5 mins) within a group. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001 using 2 way ANOVA with multiple comparisons.

Table 3 Measured blood gas variables and clinical biochemistry parameters for 12 Sprague Dawley rats (6 female, 6 male) after administration of alfaxalone at a rate of $1.67 \text{ mg kg}^{-1} \text{ min}^{-1}$ for 2.5 minutes followed by $0.75 \text{ mg kg}^{-1} \text{ min}^{-1}$ alfaxalone for 120 minutes using a one compartment infusion model. Data are mean \pm SD.

Variable	Sex	Minutes relative to the alfaxalone infusion start time				
		-5	30	60	90	120
pH *	Male	7.36 ± 0.10	7.42 ± 0.12	7.44 ± 0.08	7.41 ± 0.08	7.40 ± 0.06
	Female	7.36 ± 0.14	7.36 ± 0.14	7.35 ± 0.12	7.32 ± 0.10	7.37 ± 0.08
PCO ₂ (mmHg)	Male	36 ± 6	35 ± 12	39 ± 4	40 ± 0.4	30 ± 4
	Female	40 ± 16	44 ± 21	44 ± 16	43 ± 13	37 ± 7
PO ₂ (mmHg)	Male	140 ± 13	218 ± 19	249 ± 13	258 ± 16	276 ± 18
	Female	190 ± 31	318 ± 130	265 ± 54	263 ± 37	219 ± 70
HCO ₃ (mmol L ⁻¹)	Male	23 ± 5	25 ± 2	23 ± 3	23 ± 3	21 ± 3
	Female	24 ± 4	23 ± 4	22 ± 3	21 ± 3	21 ± 9
Sodium (mmol L ⁻¹)	Male	142 ± 2	143 ± 2	142 ± 4	143 ± 2	145 ± 3

	Female	144 ± 6	143 ± 3	147 ± 2	148 ± 5	150 ± 6
Potassium (mmol L ⁻¹)	Male	4.6 ± 0.1	4.8 ± 0.2	4.8 ± 1.1	4.7 ± 0.9	4.5 ± 2.1
	Female	5.2 ± 0.9	4.6 ± 1.2	4.8 ± 0.9	4.7 ± 1.1	5 ± 1.6
Ionized Calcium (mmol L ⁻¹)	Male	1.50 ± 0.1	1.50 ± 0.18	1.47 ± 0.24	1.37 ± 0.34	1.34 ± 0.60
	Female	1.46 ± 0.22	1.50 ± 0.13	1.51 ± 0.19	1.43 ± 0.25	1.37 ± 0.34
Chloride (mmol L ⁻¹)	Male	110 ± 3	116 ± 4	116 ± 6	114 ± 6	117 ± 5
	Female	114 ± 8	110 ± 4	116 ± 4	117 ± 7	114 ± 4
Glucose (mmol L ⁻¹)	Male	12.4 ± 3	7.1 ± 3.6	7.02 ± 3.1	5.8 ± 0.6	5.8 ± 2.8
	Female	13 ± 2.5	7.8 ± 2.5	6.3 ± 3.2	5.4 ± 1	5.9 ± 3.9
Lactate (mmol L ⁻¹)	Male	1.1 ± 0.7	1.0 ± 0.2	1.0 ± 0.1	1.1 ± 0.2	0.6 ± 0.1
	Female	1.0 ± 0.1	1.1 ± 0.4	1.0 ± 0.3	1.1 ± 0.6	0.6 ± 0.1
Creatinine (μmol L ⁻¹)	Male	20 ± 7	35 ± 16	28 ± 13	37 ± 13	33 ± 20
	Female	27 ± 12	33 ± 14	29 ± 14	37 ± 19	34 ± 14

590

591

Derived variables reported by the EPOC analyzer (actual bicarbonate, total CO₂, base excess of extra cellular fluid, base excess of blood, oxygen saturation, anion gap, anion gap potassium, haemoglobin) were calculated but are not included in the table. The baseline (-5 minute) samples were all measured under isoflurane (2% vaporizer setting) in N₂O and O₂.

* denotes statistical difference between the groups ($p < 0.05$).




Figure 1 Mean (\pm SD) alfaxalone plasma concentrations (ng mL⁻¹) for 12 Sprague Dawley rats (6 female, 6 male) after intravenous administration of alfaxalone 1.67 mg kg⁻¹ min⁻¹ for 2.5 minutes followed by 0.75 mg kg⁻¹ min⁻¹ alfaxalone for 120 minutes using a one compartment infusion model.




Figure 2 Mean (\pm SD) arterial blood pressure (MAP) (mmHg) for 12 Sprague Dawley rats (6 female, 6 male) after intravenous administration of alfaxalone $1.67 \text{ mg kg}^{-1} \text{ min}^{-1}$ for 2.5 minutes followed by $0.75 \text{ mg kg}^{-1} \text{ min}^{-1}$ alfaxalone for 120 minutes using a one compartment infusion model.