



The University of
Nottingham

UNITED KINGDOM • CHINA • MALAYSIA

Avery, Amanda and Nagar, Ravi and Hillier, Sarah and Pallister, Carolyn (2017) Impact on weight and glycaemic control in adults with diabetes attending a group- based commercial weight management programme. *Practice Nursing*, 28 (2). pp. 60-66. ISSN 0964-9271

Access from the University of Nottingham repository:

<http://eprints.nottingham.ac.uk/42453/10/Table%203.pdf>

Copyright and reuse:

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

This article is made available under the University of Nottingham End User licence and may be reused according to the conditions of the licence. For more details see:

http://eprints.nottingham.ac.uk/end_user_agreement.pdf

A note on versions:

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact eprints@nottingham.ac.uk

Table 3: Recorded weight loss and reported change in HbA1c according to length of membership at the time of survey. Data presented as means +/- standard deviations, all data was found to be normally distributed (Kolmogorov-Smirnov $p > 0.05$) and p values are for ANOVA post-hoc test with Bonferroni correction, following $p < 0.05$ ANOVA result as indicated. Significance taken at $p < 0.05$ level. As survey was open over a two week period, the average weight for members over this period was used.

	Total	< 12 weeks membership	12-24 weeks membership	24-52 weeks membership	> 52 weeks membership	p-value ANOVA	Post-Hoc
Recorded weight change (kg)	(n=393) -11.0±10.0	(n=156) -6.5±4.4	(n=65) -12.2±7.0	(n=72) -13.9±7.0	(n=99) -15.4±15.6	<0.001	Less than 12 weeks vs other groups ($p < 0.001$) No significant differences for duration of membership greater than 12 weeks
Self-reported change in HbA1c (mmol/mol) (%)	(n=156) -18 ±21 -1.6±1.9	(n=51) -18±20 -1.6±1.8	(n=24) -26±26 -2.4±2.4	(n=38) -13±23 -1.2±2.1	(n=43) -20±16 -1.8±1.5	0.126	Not Applicable