



The University of  
**Nottingham**

UNITED KINGDOM • CHINA • MALAYSIA

Avery, Amanda and Nagar, Ravi and Hillier, Sarah and Pallister, Carolyn (2017) Impact on weight and glycaemic control in adults with diabetes attending a group- based commercial weight management programme. *Practice Nursing*, 28 (2). pp. 60-66. ISSN 0964-9271

**Access from the University of Nottingham repository:**

<http://eprints.nottingham.ac.uk/42453/11/Figure%201.pdf>

**Copyright and reuse:**

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.


This article is made available under the University of Nottingham End User licence and may be reused according to the conditions of the licence. For more details see:

[http://eprints.nottingham.ac.uk/end\\_user\\_agreement.pdf](http://eprints.nottingham.ac.uk/end_user_agreement.pdf)

**A note on versions:**

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact [eprints@nottingham.ac.uk](mailto:eprints@nottingham.ac.uk)



**Figure 1:** Percentage of survey respondents achieving the NICE target for HbA1c (<48mmol/l or <6.5%) . As data is cross-sectional, it is not possible to adjust for attendance; median attendance was 6 months. n=138 for respondents with T2DM and 39 for respondents with T1DM reporting a HbA1c at time of joining, and n=128 for respondents with T2DM and 33 for respondents with T1DM at the time of survey reported a HbA1c.