

---

1

2 **WRINKLING STRUCTURES AT THE RIM OF AN INITIALLY**  
3 **STRETCHED CIRCULAR THIN PLATE SUBJECTED TO**  
4 **TRANSVERSE PRESSURE\***

5 CIPRIAN D. COMAN<sup>†</sup> AND ANDREW P. BASSOM<sup>‡</sup>

6 **Abstract.** Short-wavelength wrinkles that appear on an initially stretched thin elastic plate  
7 under transverse loading are examined. As the degree of loading is increased so wrinkles appear and  
8 their structure at the onset of buckling takes on one of three distinct forms depending on the size of  
9 the imposed stretching. With relatively little stretching, the wrinkles sit off the rim of the plate at  
10 a location which is not known a priori, but which is determined via a set of consistency conditions.  
11 These take the form of constraints on the solutions of certain coupled nonlinear differential equations  
12 that are solved numerically. As the degree of stretching grows, so an asymptotic solution of the  
13 consistency conditions is possible which heralds the structure that governs a second regime. Now the  
14 wrinkle sits next to the rim where its detailed structure can be described by the solution of suitably  
15 scaled Airy equations. In each of these first two regimes the Föppl-von Kármán bifurcation equations  
16 remain coupled, but as the initial stretching becomes yet stronger the governing equations separate.  
17 Further use of singular-perturbation arguments allows us to identify the wavelength wrinkle which  
18 is likely to be preferred in practice.

19 **Key words.** thin films, wrinkling, Föppl-von Kármán plate equations, asymptotic methods.

20 **AMS subject classifications.** 74G10, 74G60

21 **1. Introduction.** It is well known that the governing equations for thin rods,  
22 plates and shells can be obtained systematically from the general theory of nonlinear  
23 elasticity by appealing to suitable asymptotic approximations that exploit the slender-  
24 ness of such configurations. Typically, the outcome of these reduction schemes is an  
25 entire hierarchy of equations rather than a unique set; furthermore, their merit can-  
26 not be always gauged *a priori* and requires a case-by-case appraisal. The Föppl-von  
27 Kármán (FvK) nonlinear plate equations were originally derived by ad-hoc approx-  
28 imations but also represent the result of a particular asymptotic reduction (cf. [1],  
29 pp.367–447), and have proved to be a versatile choice for describing many interesting  
30 phenomena associated with thin elastic films (e.g., [2]). Arguably, this system rep-  
31 resents the simplest nonlinear model able to capture the coupling between bending  
32 deformations and the in-plane stretching of the plate mid-plane. This approximation,  
33 however, does come at a price and, despite its apparent simplicity, analytical solu-  
34 tions of the FvK system are scarce. The one notable exception is the “Euler column”  
35 [3] solution that describes a zero-Gaussian curvature deformation.

36 The principal aim of the work reported here is to throw light on a number of  
37 mathematical structures that have relevance to the FvK bifurcation system. In some  
38 recent papers [4, 5, 6] we have proposed a general asymptotic approach for describing  
39 the edge wrinkling experienced by a uniformly stretched circular elastic plate when  
40 acted upon by a transverse pressure or a concentrated central load. Generally speak-  
41 ing, the FvK bifurcation system used in our edge-wrinkling investigations is either  
42 equivalent to, or based on two coupled nonlinear equations linearised about an ax-

---

\*Submitted to the editors February 2, 2018.

<sup>†</sup>School of Mathematical Sciences, University of Nottingham, University Park, Nottingham NG7  
2RD, UK ([cdc3p@yahoo.com](mailto:cdc3p@yahoo.com))

<sup>‡</sup>School of Physical Sciences, University of Tasmania, Private Bag 37, Hobart TAS 7001, Australia  
([andrew.bassom@utas.edu.au](mailto:andrew.bassom@utas.edu.au))

43 isymmetric *nonlinear* solution that accounts for the finite mid-plane rotations in the  
 44 pre-bifurcation state.

45 The overall picture that emerged in our previous studies is summarised in Figure 1,  
 46 where typical neutral stability curves  $\mathcal{C}_\mu$  are illustrated in terms of a non-dimensional  
 47 transverse loading parameter  $\lambda > 0$  and the wrinkle mode number (or “wavenumber”)  
 48  $m \in \mathbb{N}$ . The driving system also depends on the initial in-plane stretching experienced  
 49 by the plate, which is described by a dimensionless parameter  $\mu$  defined formally in  
 50 §2 below. Strictly speaking, it is the case  $\mu \gg 1$  that lends itself most naturally to  
 51 asymptotic analysis since the energy minimum configuration for such problems – the  
 52 point  $(m_c, \lambda_c)$  in Figure 1 and corresponding to the least  $\lambda$ , can be scaled on suitable  
 53 powers of  $\mu$ . When  $0 \leq \mu \leq \mathcal{O}(1)$  the coordinates of the global minimum of  $\mathcal{C}_\mu$  are  
 54  $\mathcal{O}(1)$  quantities, a formal asymptotic strategy breaks down and critical values can  
 55 only be determined by a direct numerical simulation of the full governing equations.  
 56 In a strict mathematical sense little can be deduced for  $\mu = \mathcal{O}(1)$ , but it has been  
 57 demonstrated in [4] that even then singular perturbation methods can still be used  
 58 to provide a useful lower bound for the right-hand branch,  $\mathcal{C}_0^{(+)}$ . This is somewhat  
 59 fortuitous, but proves to be possible because, even though  $\lambda$  is formally  $\mathcal{O}(1)$ , in  
 60 practice its computed value turns out to be quite large so it can be effectively used as  
 61 an asymptotically large quantity. We remark that this is an unexpected bonus and  
 62 cannot be predicted by any formal means. Moreover, as noted in [7], the wrinkling  
 63 pattern remains strongly localised even when  $\mu = 0$  although there is no rational  
 64 theory that might suggest this could have been foreseen.




FIG. 1. The features of the neutral stability curves  $\mathcal{C}_\mu = \mathcal{C}_\mu^{(-)} \cup \mathcal{C}_\mu^{(+)}$  for the initially stretched thin elastic plate subjected to transverse pressure. The vertical axis indicates the non-dimensional pressure  $\lambda$  and the abscissa records the mode number  $m > 0$ . The parameter  $\mu \geq 0$  represents a non-dimensional measure proportional to the initial degree of radial stretching; thus, the blue curve illustrates an unstretched plate ( $\mu = 0$ ), while the red one corresponds to a taut circular configuration ( $\mu \gg 1$ ).

65 Our previous investigations have established that in the limit  $\mu \gg 1$  the FvK  
 66 system decouples and the wrinkling instability is essentially one corresponding to a  
 67 plane-stress state [8, 9], but with a nonlinear pre-buckling stress distribution. This  
 68 asymptotic decoupling plays a key role in the success of singular perturbation analyses  
 69 vis-à-vis the FvK system, a fact that is also implicit in a number of earlier works (e.g.,

70 see [10, 11]); for instance, in our problem the decoupled equations are linear and can be  
 71 solved in closed form to any order (albeit non-trivially). The question remains open as  
 72 to what happens to the FvK system as  $\mu$  is increased from zero (an unstretched plate)  
 73 to  $\mu \gg 1$  (a well-stretched plate), and it is this route to decoupling that motivates  
 74 our present study.

75 At this early stage we emphasise that our interest is with the pure buckling  
 76 problem; that is to determine the nature of  $\lambda = \lambda(\mu; m)$  which is just sufficient to  
 77 excite wrinkles of wavenumber  $m$  for a given  $\mu$ . The conventional method to isolate  
 78 the form of  $\lambda$  would be to specify  $\mu$  and then determine the corresponding  $\lambda$  as a  
 79 function of  $m$ . What makes this strategy unattractive here is that the basic solution  
 80 satisfies nonlinear equations that depend on the loading  $\lambda$ . In standard bifurcation  
 81 theory one would hope to set the basic state once and for all and then seek eigenvalues  
 82 of the perturbation equations. Here this approach will fail, or at best be complicated  
 83 to implement, as the basic equations and the system describing the wrinkles are  
 84 coupled via  $\lambda$ . The upshot would be that any critical loading values arising from the  
 85 wrinkle equations would be likely to modify the base state structure and it is unclear  
 86 how a converged solution might be arrived at which is consistent with both the base  
 87 state and wrinkle equations. Fortunately this difficulty can be neatly side-stepped  
 88 by viewing the problem from a slightly different standpoint. In this we effectively  
 89 specify  $\lambda$ , which ties down the base state, and then solve the wrinkle equations for  
 90 the wavenumber  $m$ ; it is then simple to invert the results to generate the dependence  
 91 of  $\lambda$  on  $m$ . It is a crucial feature of our work that at no stage is  $\lambda$  to be regarded as  
 92 fixed; rather for a specified  $\mu$  we are aiming to track the value of  $\lambda(m)$  just sufficient  
 93 to induce buckling. As the chosen value of  $\mu$  is changed so  $\lambda$  must compensate to  
 94 ensure we remain at the onset of buckling.

95 Within the mechanics of thin plates and shells there are several notable prece-  
 96 dents regarding the asymptotic limits of various equations as a loading parameter or a  
 97 geometrical characteristic is progressively varied. In their pioneering work [12] Junkin  
 98 and Davis studied a clamped circular plate loaded with a load on a central rigid in-  
 99 clusion by using “first-approximation” non-linear shell equations. Depending on the  
 100 magnitude of the load, they identified a sequence of plate problems that included the  
 101 usual linear equations for very small deflections and the FvK equations for moderate  
 102 deflections. A somewhat similar idea was implemented by Simmonds and Libai [13]  
 103 for a particular theory of internally pressurised spherical caps. By scaling the pressure  
 104 load and the shallowness parameters on suitable powers of a dimensionless thickness  
 105 quantity, they obtained as many as seventeen different types of simplified equations.  
 106 This suite of equations reflected a range of dominant deformation mechanisms be  
 107 they linear, nonlinear/inextensional, nonlinear/membrane or some other form. Ko-  
 108 maragiri *et al.* [14] revisited this analysis and carried out a related investigation for  
 109 a free-standing circular elastic plate under point and pressure loads. In more recent  
 110 times, Berdichevski’s asymptotic-variational technique [15] has emerged as a powerful  
 111 device that can accomplish comparable results as can be gleaned from [16] among  
 112 others. It is perhaps worth emphasising that all these studies dealt with deformation  
 113 problems, that is the load is prescribed and one tries to predict the corresponding  
 114 deformation. The problem we have in mind is somewhat different as we must tackle a  
 115 bifurcation equation. Thus, the size of the loading is intimately related to the initial  
 116 level of stretching, and can only be found by considering both the basic state and the  
 117 perturbation structure simultaneously.

118 It is acknowledged that over recent times there has been a plethora of studies  
 119 concerned with various situations in which wrinkling can arise. Researchers have

120 been concerned with developing a comprehensive framework that is able to predict  
 121 where and how tensional wrinkle patterns evolve. It appears that while many papers  
 122 deal with stretched plates, relatively few are concerned with the case when wrinkling  
 123 is provoked by an imposed transverse loading. An excellent survey of some of the  
 124 key contributions relating to plates subject to stretching or shear has been compiled  
 125 by Taylor *et al.* [17] who review the advances that have been made with geometries  
 126 such as rectangular sheets or circular regions. Our situation is somewhat different  
 127 in the sense that the wrinkling described below is generated by a transverse pressure  
 128 load. This is enough to render the basic state genuinely nonlinear and it is then not  
 129 surprising that the corresponding bifurcation equations are also distinctive. Relatively  
 130 little effort has been devoted to this class of problems although note should be made  
 131 of the numerical calculations by Adams [18] who examined the problem of a tensioned  
 132 circular plate subjected to a concentrated load.

133 The remainder of the paper is organised in the following way. We begin our  
 134 study in §2 with a quick review of the differential equations for the basic state and  
 135 the linearisation of the FvK system around this solution. A central role in our analysis  
 136 is played by a suitable large non-dimensional parameter that we shall call  $\Delta$ , and the  
 137 paper proceeds by expressing all physical quantities in terms of  $\Delta$ . In particular,  
 138 it proves possible to identify the geometry of the right-hand branch of the neutral  
 139 stability curve and trace its evolution as the *original* in-plane stretching increases.  
 140 The nonlinear axisymmetric basic state is revisited in §3 so that we can reformulate  
 141 some of the earlier features of [4] in terms of  $\Delta$ . We also show that for relatively  
 142 small  $\mu$  the associated short-wavelength wrinkle modes are governed by a parabolic  
 143 cylinder equation which is centred on a point near to, but off the rim of the plate,  
 144 and whose exact location can only be tied down upon solving a pair of consistency  
 145 conditions. These are solved numerically in §4 which shows that the structure of the  
 146 wrinkles is modified as  $\mu$  grows. Indeed, the wrinkles assume an asymptotic form,  
 147 the key elements of which are outlined in §4.1. The upshot is that a new modified  
 148 structure is appropriate to significantly enhanced  $\mu$ . At this point, which we shall  
 149 refer to as stage II, the radial extent of the wrinkles has grown but they have also  
 150 been pushed onto the rim of the plate so that an Airy-type equation becomes the  
 151 driving form. This stage II structure is developed in §5, where it is demonstrated  
 152 how a third regime must take over when  $\mu$  is enhanced further. This aspect is taken  
 153 up in §6 where it is shown how our asymptotic development automatically captures  
 154 the identity of the preferred mode when significant in-plane stretching is originally  
 155 present. The paper closes with some discussion and a few remarks.

156 **2. Formulation.** We are interested in the situation depicted in Figure 2 that  
 157 involves a circular elastic plate of uniform thickness  $h > 0$  and radius  $a$  (with  $a/h \gg$   
 158 1), a flexurally clamped edge and subjected to a uniform transverse pressure  $P$ . The  
 159 deformation of the plate is expressed using a standard cylindrical system of coordinates  
 160  $(r, \theta, z)$  defined by the usual orthonormal triad  $\{\mathbf{e}_r, \mathbf{e}_\theta, \mathbf{e}_z\}$ , with  $\mathbf{e}_z$  perpendicular to  
 161 the median plane of the plate which also contains the origin of the axes. The linearly  
 162 elastic material of the plate is characterised by the Young's modulus  $E > 0$  and the  
 163 Poisson's ratio  $0 < \nu < 1/2$ .

164 The starting point for formulating the relevant bifurcation problem is the well-  
 165 known Föppl-von Kármán (FvK) system (e.g., see [19]). When written in terms of  
 166 the transverse displacement  $w$  and a suitably defined stress function  $F$ , these become

$$167 \quad (1) \quad D\nabla^4 w - [F, w] = P \quad \text{and} \quad \nabla^4 F + \frac{Eh}{2} [w, w] = 0,$$




FIG. 2. Top and side views of a uniformly stretched circular thin plate subjected to a uniform transverse pressure; the dashed curve shown above represents its deflected shape.

168 where the first equation above accounts for the equilibrium in the normal direction,  
 169 and the second is a compatibility relation expressing the coupling between the Gaus-  
 170 sian curvature of the deformed configuration and the membrane stresses. In these  
 171 equations  $D \equiv Eh^3/12(1-\nu^2)$  represents the plate bending rigidity, and the bracket  
 172 denotes the Monge-Ampère bi-linear operator defined by  $[f, g] := (\nabla^2 f)(\nabla^2 g) - (\nabla \otimes$   
 173  $\nabla f) : (\nabla \otimes \nabla g)$  for any two smooth functions  $f$  and  $g$ . In addition  $F$  is related to  
 174 the membrane stress tensor  $\mathbf{N}$  according to  $\mathbf{N} = (\nabla^2 F)\mathbf{I}_2 - \nabla \otimes \nabla F$ , where  $\mathbf{I}_2$  is  
 175 the standard (in-plane) identity tensor  $\mathbf{I}_2 = \mathbf{e}_r \otimes \mathbf{e}_r + \mathbf{e}_\theta \otimes \mathbf{e}_\theta$ .

176 As already mentioned, the plate is clamped in the vertical direction and has  
 177 normal tractions prescribed along its circumference; this corresponds to

$$178 \quad (2a) \quad w = 0, \quad \frac{\partial w}{\partial r} = 0, \quad \text{on } r = a,$$

$$179 \quad (2b) \quad N_{rr} = N_0, \quad N_{r\theta} = 0, \quad \text{on } r = a.$$

181 To simplify (1) we set  $\rho := r/a$  and introduce the dimensionless quantities

$$182 \quad (3a) \quad \lambda := [12(1-\nu^2)]^{3/2} \left(\frac{P}{E}\right) \left(\frac{a}{h}\right)^4, \quad \mu^2 := 12(1-\nu^2) \left(\frac{N_0}{Eh}\right) \left(\frac{a}{h}\right)^2,$$

$$183 \quad (3b) \quad \bar{w} := [12(1-\nu^2)]^{1/2} \frac{w}{h}, \quad \bar{F} := 12(1-\nu^2) \frac{F}{Eh^3};$$

185 in what follows we shall drop the overbars on these re-scaled variables in order to  
 186 avoid over-complicating the notation. The parameter  $\mu^2$  measures the dimensionless  
 187 bending stiffness. In language introduced by Davidovitch *et. al* [20],  $\mu^2$  is known as

188 the bendability; it is envisaged to be fixed in an experiment while  $\lambda$  is increased until  
 189 wrinkling appears. It can then be shown that for the nonlinear axisymmetric base  
 190 state the two equations in (1) are reduced to

$$191 \quad (4) \quad \mathcal{L}_0^{(1)}[\Theta] = \lambda\rho + \frac{\Theta\Phi}{\rho} \quad \text{and} \quad \mathcal{L}_0^{(1)}[\Phi] = -\frac{\Theta^2}{2\rho},$$

192 where the new dependent variables are  $\Theta \equiv \Theta(\rho; \lambda, \mu) := dw/d\rho$  and  $\Phi \equiv \Phi(\rho; \lambda, \mu) :=$   
 193  $dF/d\rho$  with  $\mathcal{L}_0^{(k)}$  denoting the differential operator

$$194 \quad (5) \quad \mathcal{L}_0^{(k)} \equiv \frac{1}{\rho} \frac{d}{d\rho} \left( \rho \frac{d}{d\rho} \right) - \frac{k^2}{\rho^2}, \quad (k \in \mathbb{N}).$$

195 The system (4) must be solved subject to the boundary conditions

$$196 \quad (6) \quad \Theta(0) = \Theta(1) = \Phi(0) = 0, \quad \Phi(1) = \mu^2.$$

**2.1. The bifurcation boundary-value problem.** As usual, bifurcations from  
 the symmetric basic state (4) are described by a set of equations which follow easily  
 via the method of adjacent equilibrium. This involves considering perturbations to  
 the basic state  $w = \hat{w}(\rho)$ ,  $F = \hat{F}(\rho)$  which are substituted in the dimensionless  
 version of (1) and then linearized with respect to the incremental fields  $\hat{w} \equiv \hat{w}(\rho, \theta)$   
 and  $\hat{F} \equiv \hat{F}(\rho, \theta)$ . The final linear system of partial differential equations is

$$\nabla^4 \hat{w} = [\hat{w}, \hat{F}] + [\hat{w}, \hat{F}] \quad \text{and} \quad \nabla^4 \hat{F} = -[\hat{w}, \hat{w}],$$

197 which can be simplified further by looking for solutions with separable variables,

$$198 \quad (7) \quad (\hat{w}, \hat{F}) = (W(\rho), \Psi(\rho)) \cos(m\theta),$$

199 where  $m \geq 0$  is an arbitrary integer at this stage. The unknown amplitudes in (7)  
 200 satisfy the linear system

$$201 \quad (8) \quad \mathcal{L}_{11}[W] + \mathcal{L}_{12}[\Psi] = 0 \quad \text{and} \quad \mathcal{L}_{21}[W] + \mathcal{L}_{22}[\Psi] = 0,$$

202 where we have introduced the ordinary differential operators

$$203 \quad (9a) \quad \mathcal{L}_{11} \equiv [\mathcal{L}_0^{(m)}]^2 - \frac{1}{\rho} \frac{d}{d\rho} \left( \Phi \frac{d}{d\rho} \right) + \frac{d\Phi}{d\rho} \left( \frac{m}{\rho} \right)^2, \quad \mathcal{L}_{22} \equiv [\mathcal{L}_0^{(m)}]^2,$$

$$204 \quad (9b) \quad \mathcal{L}_{12} = -\mathcal{L}_{21} \equiv -\frac{1}{\rho} \frac{d}{d\rho} \left( \Theta \frac{d}{d\rho} \right) + \frac{d\Theta}{d\rho} \left( \frac{m}{\rho} \right)^2.$$

206 This eighth-order system is to be solved subject to suitable regularity conditions at  
 207 the centre of the plate together with the rim conditions (2) appropriate for a flexurally  
 208 clamped plate. In dimensionless form these constraints become simply

$$209 \quad (10) \quad W = \frac{dW}{d\rho} = \Psi = \frac{d\Psi}{d\rho} = 0, \quad \text{for } \rho \in \{0, 1\}.$$

210 Our stated intention with this work is to explore the behaviour of the FvK sys-  
 211 tem over the entire range of values of  $\mu \in [0, \infty)$  that measures the initial in-plane  
 212 stretching of the plate. Guided by our earlier remark, that even when  $\mu$  is small

213 the eigenvalue  $\lambda$  tends to be large, it is convenient to introduce the auxiliary fixed  
 214 non-dimensional parameter  $\Delta \gg 1$ . The strategy we shall adopt is to monitor the  
 215 behaviour of the system by using various quantities expressed in terms of the  $\Delta$ .  
 216 In particular results developed in [6] showed that when the wavenumber  $m$  is large  
 217 the corresponding critical loading required for wrinkling occurs at a value  $\lambda \propto m^{8/3}$ .  
 218 Guided by this we write

$$219 \quad (11) \quad \lambda = \lambda_0 \Delta^4, \quad \lambda_0 = \mathcal{O}(1),$$

220 together with the squared mode number

$$221 \quad (12) \quad m^2 = M_0 \Delta^3 + M_1 \Delta^{11/4} + M_2 \Delta^{5/2} + \dots, \quad M_j = \mathcal{O}(1), \quad (j = 0, 1, 2).$$

222 We remark that we could subsume the quantity  $M_0$  within the definition of  $\Delta$ , without  
 223 any loss of generality. However, it will prove helpful to be able to investigate various  
 224 limits while holding various physical quantities such as  $m$  or  $\mu$  fixed, and this is done in  
 225 the most transparent manner by keeping the definition of  $\Delta$  completely independent  
 226 of other quantities in the problem. Furthermore, to re-iterate the point we highlighted  
 227 in §1, although it might seem more natural to specify  $m^2$  and then seek the loading  
 228  $\lambda$  as a function of  $m$ , some mathematical subtleties inherent in the description of  
 229 the problem make this approach cumbersome. In particular, it is noted that the  
 230 basic state satisfies equations (4) which depend on  $\lambda$ . Thus if we pursue the normal  
 231 method of developing a series for  $\lambda$  as a function of  $m^2$  then there is the potential  
 232 difficulty that as we proceed we need to keep careful track of the form of the basic  
 233 state that may need to be reappraised in light of small changes to  $\lambda$ . To circumvent  
 234 this inconvenience we instead decide to determine  $m^2 := m^2(\lambda)$ . There is no formal  
 235 difficulty in adopting this viewpoint and nothing is lost so doing for once answers are  
 236 determined it is elementary to invert our results and thereby express  $\lambda := \lambda(m)$  if  
 237 preferred. At this stage there is one parameter yet to be fixed being the salient regime  
 238 for the in-plane stretching  $\mu$ , but the relevant sizing becomes evident in the course of  
 239 the calculations described below.

240 In the following we shall see that as we increase the magnitude of the dimensionless  
 241 background tension  $\mu$  the solution structures evolves through three distinct stages I–  
 242 III. Each of these is somewhat intricate and inevitably requires the introduction of  
 243 some notational complexity. Rather than minimising this by repeating symbols from  
 244 stage to stage, and thereby risking having some notation with multiple meanings in  
 245 various parts of the paper, we have chosen to have unambiguous designations. This  
 246 might initially seem overwhelming, but the three structures that are developed in  
 247 §4, §5 and §6 are separate of each other and each section can be treated as largely  
 248 self-contained. In this way, the need to undertake extensive cross-referencing between  
 249 the three calculations is hopefully mitigated as far as we are able.

250 **3. The solution structure for  $\Delta \gg 1$ : stage I.** Given the form of (11),  
 251 simple scaling arguments applied to the base-state equations (4) suggest that across  
 252 the majority of circular plate, where  $\rho = \mathcal{O}(1)$ , we have

$$253 \quad (13) \quad \Theta = \Delta^{4/3} \Theta_0 + \Delta^{-4/3} \Theta_1 \dots, \quad \Phi = \Delta^{8/3} \Phi_0 + \Phi_1 + \dots$$

254 Leading-order terms in (4) reduce to

$$255 \quad (14) \quad \Theta_0 \Phi_0 = -\lambda_0 \rho^2, \quad \mathcal{L}_0^{(1)}[\Phi_0] = -\frac{\Theta_0^2}{2\rho},$$



256 from which it quickly follows that

$$257 \quad (15) \quad \mathcal{L}_0^{(1)}[\Phi_0] = -\frac{\lambda_0^2 \rho^3}{2\Phi_0^2}.$$

258 It is a routine exercise to show that at the centre of the plate  $\Phi_0 \sim A\rho + \mathcal{O}(\rho^3)$   
 259 for some constant  $A \in \mathbb{R}$  that could be determined numerically, but whose value is  
 260 immaterial for our immediate purposes. Rather, what is of more significance is the  
 261 nature of the solution of (14) at the rim  $\rho \rightarrow 1^-$ . In view of the boundary conditions  
 262 (6) on the base state at  $\rho = 1$  we anticipate that, if  $\mu$  is small, then  $\Phi_0 \rightarrow 0$  as  $\rho \rightarrow 1^-$ ,  
 263 which requires

$$264 \quad (16) \quad \Phi_0 \sim \delta \left( x^{2/3} + \dots \right), \quad \Theta_0 \sim -\frac{\lambda_0}{\delta} x^{-2/3} (1 - \dots), \quad \text{where } \delta \equiv \left( \frac{3}{2} \lambda_0 \right)^{2/3},$$

265 as  $x \equiv 1 - \rho \rightarrow 0^+$ . (This expression follows immediately from enforcing the balance  
 266 between the second derivative on the left hand side of (15) with the nonlinear term  
 267 on the right hand side.)

268 This then highlights the significance of a suitable rim layer wherein the majority of  
 269 the wrinkling will take place. Elementary scaling of the governing equations suggests  
 270 that  $x = \mathcal{O}(\Delta^{-1})$ , so we define

$$271 \quad (17) \quad \rho = 1 - \frac{X}{\Delta}, \quad X = \mathcal{O}(1)$$

272 whereupon, governed by the behaviours (16), we expect that

$$273 \quad (18) \quad (\Phi, \Theta) = \Delta^2(\phi_0, \theta_0) + \Delta^{4/3}(\phi_1, \theta_1) + \dots$$

274 If a dash denotes differentiation with respect to  $X$ , then substitution in (4) shows  
 275 that the zeroth-order terms satisfy

$$276 \quad (19) \quad \theta_0'' = \lambda_0 + \phi_0 \theta_0, \quad \phi_0'' = -\frac{1}{2} \theta_0^2,$$

277 and matching with the outer behaviour (16) demands that  $\phi_0 \sim \delta X^{2/3}$  and  $\theta_0 \sim$   
 278  $-(\lambda_0/\delta)X^{-2/3}$  as  $X \rightarrow \infty$ .

279 It is the rim condition  $\Phi(1) = \mu^2$  from (6) that provides the clue for the appro-  
 280 priate scaling for  $\mu$ . If we put

$$281 \quad (20) \quad \mu = \Delta \mu_0, \quad \mu_0 = \mathcal{O}(1),$$

282 then we must have

$$283 \quad (21) \quad \theta_0(0) = 0 \quad \text{and} \quad \phi_0(0) = \mu_0^2.$$

284 Clearly, the value of  $\mu_0 > 0$  plays a significant role in setting the leading-order form  
 285 of the basic state within the rim region and thus, presumably, is important in setting  
 286 the loading that generates wrinkle modes. Hence we now work with  $\mu_0$  assumed fixed  
 287 and given, and seek to determine the value of  $\lambda_0(m)$  that marks the onset of buckling.



288 **3.1. The structure of the eigen-deformation.** Given the form of the basic  
 289 solution we are able to proceed to examine the perturbation equations. We can  
 290 thereby identify the key scalings that ensure that quadratic terms drive perturbations  
 291 that are confined to a thin zone within the  $X = \mathcal{O}(1)$  region. Put another way, the  
 292 basic state develops a rim layer and inside that, already thin, layer sit the wrinkle  
 293 modes. It can be verified that this inner rim later is of relative extent  $\mathcal{O}(\Delta^{-1/4})$ , and  
 294 so the wrinkle exists at some location  $X = X_0$  about which we define the rescaled  
 295 variable  $Y = \mathcal{O}(1)$ ,

$$296 \quad (22) \quad Y := \Delta^{1/4}(X - X_0) \quad \text{or} \quad \rho = 1 - \frac{X_0}{\Delta} - \frac{Y}{\Delta^{5/4}}.$$

297 There are now two issues to be settled: (i) what sets the value of the location  $X_0$  and  
 298 (ii) how is the disturbance confined to the vicinity of this point? We can begin to  
 299 address these questions by expanding the rim layer quantities  $\phi_j(X)$  as Taylor series  
 300 taken about  $X = X_0$ . This shows that where  $Y = \mathcal{O}(1)$  we have

$$302 \quad (23) \quad \Phi = \Delta^2 \left( \phi_{00} + \Delta^{-1/4} \phi_{01} Y + \frac{1}{2} \Delta^{-1/2} \phi_{02} Y^2 + \frac{1}{6} \Delta^{-3/4} \phi_{03} Y^3 + \dots \right)$$

$$303 \quad \quad \quad + \Delta^{4/3} \left( \phi_{10} + \Delta^{-1/4} \phi_{11} Y + \dots \right),$$

305 where the constants  $\phi_{ij}$  denote the  $j^{\text{th}}$  derivative of  $\phi_i(X)$  evaluated at  $X = X_0$ .  
 306 Taking derivatives shows that

$$307 \quad (24) \quad \frac{d\Phi}{d\rho} = -\Delta^3 \phi_{01} - \Delta^{11/4} \phi_{02} Y - \frac{1}{2} \Delta^{5/2} \phi_{03} Y^2 + \dots$$

308 and we remark that expressions completely analogous to (23) and (24) hold for  $\Theta$  and  
 309 its derivative, with the  $\phi_{ij}$  replaced by  $\theta_{ij}$  which represents the  $j^{\text{th}}$ -order derivative  
 310 of  $\theta_i(X)$  evaluated at  $X = X_0$ . Notice that although the base state correction term  
 311  $\phi_1$  enters both the expressions (18) and the Taylor series (23), it is not required for  
 312 the results we derive below. Hence, for reasons of brevity, we do not discuss  $\phi_1$  (and  
 313  $\theta_1$ ) further here, though of course their presence would have to be properly accounted  
 314 for if we were to delve deeper into later terms in our series solutions.

315 **3.2. The bifurcation equations.** Given these proposed structures, and with  
 316 the squared mode number  $m^2$  defined by (12), the scene is now set for determining  
 317 the important equations. We look for a solution of (8) of the form

$$318 \quad (25) \quad (W, \Psi) = (W_0, \Psi_0) + \Delta^{-1/4}(W_1, \Psi_1) + \Delta^{-1/2}(W_2, \Psi_2) + \dots,$$

319 and remember that  $\Phi$  and its derivative are given by (23) and (24). On substituting  
 320 (25) into the original equations (8), collecting like powers of  $\Delta$ , and then setting to  
 321 zero their corresponding coefficients results in a hierarchy of coupled equations, as  
 322 explained below.

323 Terms of  $\mathcal{O}(\Delta^6)$  in the two equations yield

$$324 \quad (26a) \quad \mathcal{R}_1[W_0, \Psi_0] \equiv (M_0 - \phi_{01})W_0 - \theta_{01}\Psi_0 = 0,$$

$$325 \quad (26b) \quad \mathcal{R}_2[W_0, \Psi_0] \equiv \theta_{01}W_0 + M_0\Psi_0 = 0.$$

327 The consistency of this linear homogeneous system in  $W_0$  and  $\Psi_0$  requires

$$328 \quad (27) \quad M_0(M_0 - \phi_{01}) + \theta_{01}^2 = 0.$$

329 At  $\mathcal{O}(\Delta^{23/4})$  it follows that

$$330 \quad (28a) \quad \mathcal{R}_1[W_1, \Phi_1] = Y\theta_{02}\Psi_0 - (M_1 - \phi_{02}Y)W_0,$$

$$331 \quad (28b) \quad \mathcal{R}_2[W_1, \Phi_1] = -\theta_{02}YW_0 - M_1\Psi_0.$$

333 Again, a solution is only possible if suitable consistency conditions hold. The pair of  
334 equations (26) imply that  $M_0\mathcal{R}_1(W, \Phi) + \theta_{01}\mathcal{R}_2(W, \Phi) \equiv 0$  as these two operators are  
335 linearly related. It follows that the system (28) is compatible only if

$$336 \quad (29) \quad M_1 = 0 \quad \text{and} \quad M_0\phi_{02} = 2\theta_{01}\theta_{02}.$$

337 We need to proceed as far as  $\mathcal{O}(\Delta^{11/2})$ . We determine that

$$338 \quad \mathcal{R}_1[W_2, \Phi_2] = \theta_{02}Y\Psi_1 + \phi_{02}YW_1 + \frac{1}{2}\theta_{03}Y^2\Psi_0 - \left(M_2 - \frac{1}{2}\phi_{03}Y^2\right)W_0 + 2\frac{d^2W_0}{dY^2},$$

$$339 \quad \mathcal{R}_2[W_2, \Phi_2] = -\theta_{02}YW_1 - \frac{1}{2}\theta_{03}Y^2W_0 - M_2\Psi_0 + 2\frac{d^2\Psi_0}{dY^2}.$$

341 The consistency of this pair requires

$$342 \quad (31) \quad \frac{d^2W_0}{dY^2} + \left[\frac{M_0\phi_{03} - 2\theta_{01}\theta_{03} - 2\theta_{02}^2}{4(2M_0 - \phi_{01})}\right]Y^2W_0 - \frac{1}{2}M_2W_0 = 0,$$

343 which, when cast in the generic form

$$344 \quad (32) \quad \frac{d^2W_0}{dY^2} - \gamma Y^2W_0 + \delta W_0 = 0,$$

345 admits the exact solution  $W_0 \propto \exp(-\gamma^{1/2}Y^2/2)$  if  $\delta = \gamma^{1/2}$ . This gives

$$346 \quad (33) \quad M_2 = -2 \left[\frac{2\theta_{02}^2 + 2\theta_{01}\theta_{03} - M_0\phi_{03}}{4(2M_0 - \phi_{01})}\right]^{1/2},$$

347 as long as  $\gamma > 0$ . The expression  $W_0(Y) \propto \exp(-\gamma^{1/2}Y^2/2)$  proves that the solution  
348 is effectively confined to the  $Y = \mathcal{O}(1)$  region subsumed within the  $X = \mathcal{O}(1)$  rim  
349 layer governing the base structure.

350 We now have the information we require to uncover the location of the wrinkles  
351 centred at  $X = X_0$ . For a given  $\lambda_0$  the leading-order rim solution  $(\phi_0, \theta_0)$  satisfies  
352 the coupled system (19) subject to (21) and the matching conditions  $\phi_0 \sim \delta X^{2/3}$   
353 and  $\theta_0 \sim -(\lambda_0/\delta)X^{-2/3}$  as  $X \rightarrow \infty$ . Wrinkling occurs with a scaled square mode  
354 number  $M_0$  and is located at  $X = X_0$ , where  $M_0$  and  $X_0$  are determined by solving the  
355 consistency equations (27) and (29). Solution of this problem requires some associated  
356 numerical work, as explained briefly in the next section.

357 **4. Numerical solution of the stage-I equations.** Our computational task  
358 requires that, *given* the scaled constant  $\mu_0 > 0$ , we need to determine the relationship  
359 between  $\lambda_0$  and  $M_0$ . It turns out that considerable simplification can be achieved by  
360 some judicious scaling. If we define a new rim co-ordinate  $\hat{X} \geq 0$  according to

$$361 \quad (34) \quad \hat{X} := \lambda_0^{1/4}X,$$

362 and write the base structure variables  $\phi_0 =: \lambda_0^{1/2} \widehat{\phi}_0$  and  $\theta_0 =: \lambda_0^{1/2} \widehat{\theta}_0$ , then it follows  
 363 that

$$364 \quad (35) \quad \widehat{\theta}'_0 = 1 + \widehat{\phi}_0 \widehat{\theta}_0, \quad \widehat{\phi}'_0 = -\frac{1}{2} \widehat{\theta}_0^2,$$

365 subject to the constraints

$$366 \quad (36) \quad \widehat{\phi}_0 \sim \alpha \widehat{X}^{2/3} + \dots, \quad \widehat{\theta}_0 \sim -\frac{1}{\alpha} \widehat{X}^{-2/3} + \dots \quad \text{as } \widehat{X} \rightarrow \infty; \quad \alpha \equiv \left(\frac{3}{2}\right)^{2/3}$$

367 together with

$$368 \quad (37) \quad \widehat{\theta}_0(0) = 0 \quad \text{and} \quad \widehat{\phi}_0(0) = \widehat{\Lambda};$$

369 here, we have introduced the definition

$$370 \quad (38) \quad \widehat{\Lambda} := \frac{\mu_0^2}{\lambda_0^{1/2}}.$$

371 If furthermore, we put  $M_0 =: \lambda_0^{3/4} \widehat{M}_0$  and denote by  $\widehat{\phi}_{0j}$  and  $\widehat{\theta}_{0j}$  the  $j^{\text{th}}$  derivatives  
 372 of  $\widehat{\phi}_0$  and  $\widehat{\theta}_0$  evaluated at  $\widehat{X} = \widehat{X}_0$ , then the consistency conditions (27) and (29)  
 373 become just

$$374 \quad (39) \quad \widehat{M}_0(\widehat{M}_0 - \widehat{\phi}_{01}) + \widehat{\theta}_{01}^2 = 0 \quad \text{and} \quad \widehat{M}_0 \widehat{\phi}_{02} = 2\widehat{\theta}_{01} \widehat{\theta}_{02}.$$

375 By this device we have reduced by one the dimension of the parameter space over  
 376 which solution is required. For each  $\widehat{\Lambda}$  there is one pair of corresponding  $(\widehat{M}_0, \widehat{X}_0)$  and  
 377 we are faced with a three-point boundary-value problem comprising the fourth-order  
 378 system (35)–(37) subject to consistency conditions to be imposed at a point  $\widehat{X}_0$  that  
 379 is part of the solution. This computation was carried out using standard routines  
 380 available in MATLAB.

381 Some representative solutions are shown in Figure 3; in the left panel is illustrated  
 382 the dependence of  $\widehat{M}_0$  on  $\widehat{\Lambda}$ , while the right panel indicates the corresponding form of  
 383 the location  $\widehat{X}_0$  within the rim region. We note that for no initial in-plane stretching,  
 384 i.e.  $\widehat{\Lambda} = 0$ , we have finite values  $\widehat{M}_0 \simeq 0.8721$  and  $\widehat{X}_0 \simeq 1.066$ . As  $\widehat{\Lambda}$  increases  
 385 so initially  $\widehat{X}_0$  grows, but this trend is soon reversed and both  $\widehat{M}_0$  and  $\widehat{X}_0$  drop  
 386 steadily with  $\widehat{\Lambda}$ . This suggests that to account for stronger stretching  $\mu_0 \gg 1$  (and  
 387 so  $\widehat{\Lambda} \gg 1$  by definition (38)) some sort of new structure ought to come into play in  
 388 an appropriate large- $\widehat{\Lambda}$  limit. To unravel the corresponding details the first step is  
 389 therefore to examine the nature of the solution of (35)–(37) subject to (39) as  $\widehat{\Lambda} \rightarrow \infty$ .

390 **4.1. The solution of (35)–(39) for large  $\widehat{\Lambda}$ .** Consideration of the boundary  
 391 condition imposed on  $\widehat{\phi}_0$  at  $\widehat{X} = 0$  together with the nature of the governing equations  
 392 suggest that when  $\widehat{\Lambda} \gg 1$  the solution develops a short-scale structure on a length  
 393  $\mathcal{O}(\widehat{\Lambda}^{-1/2})$ . We therefore define

$$394 \quad (40) \quad \widehat{X} = \widehat{\Lambda}^{-1/2} z,$$

395 and propose that the solution takes the form

$$396 \quad (41) \quad \widehat{\theta}_0 = \widehat{\Lambda}^{-1} \widetilde{\theta}_0(z) + \widehat{\Lambda}^{-3} \widetilde{\theta}_1(z) + \dots, \quad \widehat{\phi}_0 = \widehat{\Lambda} \widetilde{\phi}_0(z) + \widehat{\Lambda}^{-1} \widetilde{\phi}_1(z) + \widehat{\Lambda}^{-3} \widetilde{\phi}_2(z) + \dots$$




FIG. 3. The forms of  $\widehat{M}_0$  (left) and  $\widehat{X}_0$  (right) as functions of the parameter  $\widehat{\Lambda}$  as determined from the solution of system (35)–(37) subject to (39) being satisfied at  $\widehat{X} = \widehat{X}_0$ . Shown as superimposed markers are the corresponding large- $\widehat{\Lambda}$  asymptotic results (46).

397 On substituting these forms in (35), comparison of like coefficients of  $\widehat{\Lambda}$  in the two  
398 equations yield that

$$399 \quad \frac{d^2 \tilde{\theta}_0}{dz^2} = 1 + \tilde{\theta}_0 \tilde{\phi}_0, \quad \frac{d^2 \tilde{\theta}_1}{dz^2} = \tilde{\theta}_0 \tilde{\phi}_1 + \tilde{\theta}_1 \tilde{\phi}_0, \quad \frac{d^2 \tilde{\phi}_0}{dz^2} = \frac{d^2 \tilde{\phi}_1}{dz^2} = 0 \quad \text{and} \quad \frac{d^2 \tilde{\phi}_2}{dz^2} = -\frac{1}{2} \tilde{\theta}_0^2.$$

400 In view of the boundary conditions we suppose that  $\tilde{\phi}_0 \equiv 1$ , a claim that can be  
401 checked later. Given this, it follows quickly that  $\tilde{\theta}_0 = -1 + \exp(-z)$  and we can also  
402 deduce that  $\tilde{\phi}_1$  is proportional to  $z$ . We cannot tie down this solution completely  
403 without recourse to the far-field conditions (36) for  $\widehat{X} \rightarrow \infty$ . The fact that the  
404 solution does not match directly onto the far-field requirements suggests strongly  
405 that the inner-solution zone must be supplemented by some form of outer structure.  
406 It is not difficult to verify that this outer zone lies where  $\widehat{X} = \widehat{\Lambda}^{3/2} \tilde{Y}$  with  $\tilde{Y} = \mathcal{O}(1)$   
407 and that here

$$408 \quad (42) \quad \tilde{\theta}_0 = \widehat{\Lambda}^{-1} \tilde{\Theta}_0(\tilde{Y}) + \dots, \quad \tilde{\phi}_0 = \widehat{\Lambda} \tilde{\Phi}_0(\tilde{Y}) + \dots$$

To match with the inner region requires that  $\tilde{\Theta}_0 \rightarrow -1$  and  $\tilde{\Phi}_0 \rightarrow 1$  as  $\tilde{Y} \rightarrow 0$ .  
Leading-order terms in the governing equations (35) give

$$\tilde{\Theta}_0 \tilde{\Phi}_0 + 1 = 0 \quad \text{and} \quad \frac{d^2 \tilde{\Phi}_0}{d\tilde{Y}^2} = -\frac{1}{2} \tilde{\Theta}_0^2 \quad \implies \quad \frac{d^2 \tilde{\Phi}_0}{d\tilde{Y}^2} = -\frac{1}{2 \tilde{\Phi}_0^2}.$$

409 This latter equation admits the exact solution  $\tilde{\Phi}_0 = (1 + 3\tilde{Y}/2)^{2/3}$  which hence yields  
410 that  $\tilde{\Theta}_0 = -(1 + 3\tilde{Y}/2)^{-2/3}$ . These expressions match automatically with the far-field  
411 requirements (36) and with the inner-zone solutions as  $\tilde{Y} \rightarrow 0$ . Furthermore we can  
412 now deduce that  $\tilde{\phi}_1 = z$ , rather than just being proportional to it, and it is then a

413 routine matter to solve for  $\tilde{\phi}_2$ . Taken together, this means that within the inner zone  
414 (40)

$$415 \quad (43a) \quad \hat{\theta}_0 = \hat{\Lambda}^{-1}[-1 + \exp(-z)] + \dots,$$

$$416 \quad (43b) \quad \hat{\phi}_0 = \hat{\Lambda} + \hat{\Lambda}^{-1}z + \hat{\Lambda}^{-3} \left[ -\frac{1}{4}z^2 + \exp(-z) - \frac{1}{8}\exp(-2z) + \tilde{c}z - \frac{7}{8} \right] + \dots,$$

417

418 where the precise value of  $\tilde{c} \in \mathbb{R}$  will not be required.

419 We have now shown that when  $\hat{\Lambda} \gg 1$  the appropriate solution of (35)–(37)  
420 develops a two-layer structure with an inner  $\mathcal{O}(\hat{\Lambda}^{-1/2})$ -zone and a wider outer region.  
421 We still need to identify the corresponding values of  $\hat{M}_0$  and  $\hat{X}_0$  that together fulfil the  
422 consistency requirements (39). The numerical solutions sketched in Figure 3 suggested  
423 that as  $\hat{\Lambda} \rightarrow \infty$  so  $\hat{X}_0 \rightarrow 0$ , and therefore it is unlikely that the consistency conditions  
424 will hold somewhere in the outer zone. Granted this, suppose that (39) apply at some  
425 point  $\hat{X}_0 = \hat{\Lambda}^{-1/2}z_0$  for some  $z_0 > 0$  to be found. In order to satisfy (39) it is clear  
426 that the values of the various derivatives  $\hat{\theta}_{01}$ ,  $\hat{\theta}_{02}$ ,  $\hat{\phi}_{01}$  and  $\hat{\phi}_{02}$  need to be found. These  
427 can be inferred directly from our foregoing results (43), which yield

$$428 \quad (44) \quad \hat{\phi}_{01} = \hat{\Lambda}^{-1/2} + \dots, \quad \hat{\theta}_{01} = -\hat{\Lambda}^{-1/2} \exp(-z_0) + \dots,$$

$$429 \quad (45) \quad \hat{\phi}_{02} = -\frac{1}{2}\hat{\Lambda}^{-2}(1 - \exp(-z_0))^2 + \dots, \quad \hat{\theta}_{02} = \exp(-z_0) + \dots$$

430

431 We proceed by examining the first of the consistency conditions in (39). Given  
432 the values noted in (44) there appear to be two possibilities: either  $\hat{M}_0 \simeq -\hat{\theta}_{01}^2/\hat{\phi}_{01}$   
433 or  $\hat{M}_0 \simeq \hat{\phi}_{01} = \hat{\Lambda}^{-1/2}$ . If we take the former option, routine algebra shows that the  
434 second consistency requirement  $\hat{M}_0\hat{\phi}_{02} = 2\hat{\theta}_{01}\hat{\theta}_{02}$  cannot be satisfied. We are then  
435 left to conclude that

$$436 \quad (46) \quad \hat{M}_0 \simeq \hat{\Lambda}^{-1/2} \quad \text{and} \quad \hat{X}_0 \simeq \hat{\Lambda}^{-1/2} \ln(2\hat{\Lambda}), \quad \text{as} \quad \hat{\Lambda} \rightarrow \infty,$$

437 where the value of  $\hat{X}_0$  follows from the second consistency condition. These large-  
438  $\hat{\Lambda}$  predictions are superimposed on the results shown in Figure 3, and both show  
439 excellent agreement with the direct numerical simulations.

440 **5. The emergence of structure for larger values of  $\mu$ : stage II.** In  
441 the preceding sections we have sought to explain the structure of wrinkling eigen-  
442 deformations with high mode numbers ( $m \sim \Delta^{3/2}$ ) when the in-plane stretching  
443 parameter  $\mu$  is of size  $\mathcal{O}(\Delta)$ : recall (12) and (20). Equation (11) reminds us that  
444 the corresponding loading for wrinkling is  $\mathcal{O}(\Delta^4)$  and we now investigate how the  
445 situation needs to be modified as  $\underline{\mu}$  grows. The workings of the previous section  
446 show what is likely to happen as  $\hat{\Lambda}$  increases. In particular we observe that the  
447 square of the mode number  $m^2 \sim \Delta^3 M_0 = \Delta^3 \lambda_0^{3/4} \hat{M}_0$ , and for  $\hat{\Lambda} \gg 1$  we predicted  
448  $\hat{M}_0 \sim \hat{\Lambda}^{-1/2} = \lambda_0^{1/4}/\mu_0$  by definition (38). Hence the wrinkle mode number is

$$449 \quad (47) \quad m \sim \Delta^{3/2} \lambda_0^{1/2} / \mu_0^{1/2}.$$

We need to be careful that we continue to examine eigenstates with mode numbers  
consistent with those appropriate to stage I; that is, we should keep  $m \sim \Delta^{3/2}$ . This  
then suggests  $\lambda_0 \sim \mu_0$  while a second constraint for fixing the appropriate sizings for

$\lambda_0$  and  $\mu_0$  follows from the wrinkling structure itself. Using equation (32) and the scalings of §3, it follows that the wrinkling layer is of extent  $\mathcal{O}(\Delta^{-5/4}\widehat{\gamma}^{-1/4})$ , where

$$\widehat{\gamma} \equiv \lambda_0^{5/4} \left[ \frac{-\widehat{M}_0\widehat{\phi}_{03} + 2\widehat{\theta}_{01}\widehat{\theta}_{03} + 2\widehat{\theta}_{02}^2}{4(2\widehat{M}_0 - \widehat{\phi}_{01})} \right].$$

450 Given the asymptotic results (44), it transpires that the depth of the wrinkle zone is  
451 comparable to the distance of its centre from the rim when

$$452 \quad (48) \quad \Delta^{-5/4}\lambda_0^{-1/4}\mu_0^{-1/4} \sim \Delta^{-1} \quad \text{or} \quad \lambda_0\mu_0 \sim \Delta.$$

453 Taken with our earlier expectation  $\lambda_0 \sim \mu_0$  it is now clear that some new structure is  
454 anticipated once  $\mu \equiv \Delta\mu_0$  becomes  $\mathcal{O}(\Delta^{3/2})$ . To avoid introducing a plethora of new  
455 variables we recycle much of the preceding notation, changing only those parts that  
456 are crucial to avoid confusion.

457 Our discussion immediately above suggests that the stretching and loading must  
458 be scaled according to

$$459 \quad (49) \quad \mu = \Delta^{3/2}\mu_0^\dagger, \quad \lambda = \Delta^{9/2}\lambda_0^\dagger, \quad \text{for some } \mu_0^\dagger, \lambda_0^\dagger = \mathcal{O}(1),$$

460 which replace (20) and (11), respectively. We have been careful to ensure that we  
461 continue to seek eigen-deformations with mode numbers  $\mathcal{O}(\Delta^{3/2})$ , so suppose that

$$462 \quad (50) \quad m^2 = M_0^\dagger\Delta^3 + M_1^\dagger\Delta^2 + \dots$$

463 Once again, we proceed assuming that  $\mu_0^\dagger$  is *fixed* and given, and endeavour to find  
464 the form of  $\lambda_0^\dagger = \lambda_0^\dagger(M_0^\dagger)$  that marks the onset of buckling. Our previous asymptotics  
465 predict that the wrinkling is confined to an  $\mathcal{O}(\Delta^{-1})$ -distance off the rim, so we can  
466 simply retain definition (17) with  $\rho = 1 - X\Delta^{-1}$ .

467 In view of the increase in the loading  $\lambda$  the basic state is modified, though the key  
468 equations are only slightly altered. The basic state across the majority of the plate  
469 now satisfies

$$470 \quad (51) \quad \Theta = \Delta^{3/2}\Theta_0 + \dots, \quad \Phi = \Delta^3\Phi_0 + \dots,$$

471 where

$$472 \quad (52) \quad \Theta_0\Phi_0 = -\lambda_0^\dagger\rho^2 \quad \text{and} \quad \mathcal{L}_0^{(1)}[\Phi_0] = -\frac{(\lambda_0^\dagger)^2\rho^3}{2\Phi_0^2}.$$

473 Previously we needed to solve for  $\Phi_0$  subject to the requirement that it vanished as  
474  $\rho \rightarrow 1^-$ ; however, now the enhanced value of  $\mu$  in (49) means that we simply require  
475 that  $\Phi_0(1) = (\mu_0^\dagger)^2$ . If we write  $\Phi_0 \equiv (\mu_0^\dagger)^2\phi_0$  then it follows that

$$476 \quad (53) \quad \mathcal{L}_0^{(1)}[\phi_0] = -\frac{\Gamma^2\rho^3}{2(\phi_0)^2}; \quad \phi_0(0) = 0, \quad \phi_0(1) = 1; \quad \Gamma \equiv \frac{\lambda_0^\dagger}{(\mu_0^\dagger)^3}.$$

477 We need to ascertain the behaviour of this solution in the rim zone  $X = \mathcal{O}(1)$  and it  
478 is straightforward to deduce that if  $\phi_0'(1) \equiv \beta$  then in the rim zone

$$479 \quad (54) \quad \Phi = \Delta^3(\mu_0^\dagger)^2 \left[ 1 - \frac{\beta X}{\Delta} + \frac{1}{2} \left( 1 - \beta - \frac{1}{2}\Gamma^2 \right) \frac{X^2}{\Delta^2} + \dots \right]; \quad \Theta = \mathcal{O}(\Delta^{3/2}).$$

480 If where  $X = \mathcal{O}(1)$  the wrinkle adopts the form

481 (55) 
$$(W, \Psi) = (W_0^\dagger, \Psi_0^\dagger) + \Delta^{-1}(W_1^\dagger, \Psi_1^\dagger) + \dots,$$

482 then leading-order terms arising from substitution in (8) tell us that

483 (56) 
$$M_0^\dagger + \beta(\mu_0^\dagger)^2 = 0 \quad \text{and} \quad (\mu_0^\dagger)^2 M_0^\dagger \Psi_0^\dagger = \lambda_0^\dagger(\beta - 2)W_0^\dagger.$$

484 At next order in equation (8a) we find that

485 (57) 
$$\left[ 2 + \frac{(\mu_0^\dagger)^2}{M_0^\dagger} \right] \frac{d^2 W_0^\dagger}{dX^2} - \left[ (\mu_0^\dagger)^2 \left( \beta + \frac{1}{2}\Gamma^2 - 1 \right) X + M_1^\dagger \right] W_0^\dagger = 0.$$

486 This equation is merely a scaled form of the ubiquitous Airy equation  $y'' - xy = 0$ ,  
 487 which is known to admit a solution with  $y(x_0) = 0$  and  $y \rightarrow 0$  as  $x \rightarrow \infty$  if  $x_0 \simeq$   
 488  $-2.331$ . Given this, we deduce that equation (57) enables  $W_0^\dagger \rightarrow 0$  both as  $X \rightarrow 0$   
 489 and as  $X \rightarrow \infty$  if

490 (58) 
$$M_1^\dagger \simeq -2.331(M_0^\dagger)^{-1/3} \left[ \left( \beta + \frac{1}{2}\Gamma^2 - 1 \right) (\mu_0^\dagger)^2 + 2M_0^\dagger \right]^{2/3} \left[ 2M_0^\dagger + (\mu_0^\dagger)^2 \right]^{1/3}.$$

491 We now have the elements required to determine the loading parameter  $\lambda_0^\dagger$  in  
 492 terms of  $\mu_0^\dagger$ . The key to unlocking this dependence lies in the requirement  $\phi_0'(1) \equiv \beta$   
 493 and the consistency condition (56a) combined with the basic state equation (53).  
 494 This second-order equation already is subject to the two requirements,  $\phi_0(0) = 0$  and  
 495  $\phi_0(1) = 1$ , and the third constraint  $\phi_0'(1) = -M_0^\dagger/(\mu_0^\dagger)^2$ , which follows directly from  
 496  $\phi_0'(1) \equiv \beta$  and (56a), means that a solution only exists for certain values of  $\Gamma$ . We  
 497 can write this in the alternative form

498 (59) 
$$\lambda_0^\dagger = (\mu_0^\dagger)^3 G \left[ \frac{M_0^\dagger}{(\mu_0^\dagger)^2} \right],$$

499 for some function  $G[\cdot]$  that can only be determined numerically; the form of this  
 500 function is illustrated in Figure 4.

501 It is a straightforward computational exercise to show that problem (53) admits  
 502 a solution with  $\phi_0'(1) = 0$  when  $\Gamma = K_0 \approx 3.212$ . This then tells us that for large  $\mu_0^\dagger$   
 503 (and small  $\beta$ ) then  $\lambda_0^\dagger \simeq K_0(\mu_0^\dagger)^3$ . Moreover, if we look for a solution of (53) as a  
 504 regular series in inverse powers of  $\mu_0^\dagger$  we can derive the two-term result

505 (60) 
$$\lambda_0^\dagger = K_0(\mu_0^\dagger)^3 + 1.217M_0^\dagger\mu_0^\dagger + \dots,$$

506 which is included on Figure 4. It is observed that agreement is excellent, even for  
 507 surprisingly modest values of  $\mu_0^\dagger$ .

508 These features forecast the expected behaviours at even larger values of  $\mu$ . As  $\mu_0^\dagger$   
 509 grows so the leading order loading parameter  $\lambda_0^\dagger$  becomes independent of the mode  
 510 number  $M_0^\dagger$ , and the fact that the quantity  $M_1^\dagger \sim \mathcal{O}((\mu_0^\dagger)^2)$ , according to (58), means  
 511 that a restructuring should be anticipated once  $\mu_0^\dagger = \mathcal{O}(\Delta^{1/2})$ . Then  $\mu = \mathcal{O}(\Delta^2)$  and  
 512 this last stage is described next.






FIG. 4. The dependence of  $\lambda_0^\dagger$  on  $\mu_0^\dagger$  according to (59) when  $M_0^\dagger = 1$ . Shown superimposed on this plot are the one- and two-term asymptotic results (60), which correspond to the triangular and round markers, respectively.

513 **6. Stage III: strong stretching.** Guided by the previous analysis we can  
 514 quickly sketch the structure appropriate when

$$515 \quad (61) \quad \mu = \Delta^2 \tilde{\mu}, \quad \text{with} \quad \tilde{\mu} = \mathcal{O}(1).$$

516 We anticipate that once more  $m^2 = \Delta^3 \tilde{M}$ , for some  $\tilde{M} = \mathcal{O}(1)$ , but that the com-  
 517 mensurate loading is now

$$518 \quad (62) \quad \lambda = K_0 \tilde{\mu}^3 \Delta^6 + \tilde{\lambda}_1 \Delta^5 + \tilde{\lambda}_2 \Delta^4 + \dots,$$

where  $\lambda_j = \mathcal{O}(1)$  ( $j = 1, 2$ ) are yet to be determined. In passing we remark that this scaling that  $m \sim \mu^{3/4}$  was first derived using asymptotic arguments by Coman & Bassom [8] and later Davidovitch *et al.* [20] gave a simple argument based on scaling to confirm this result. Earlier in the paper we stressed our desire to take a solitary one-term form for  $\lambda$  and, at face value, it seems that we are now deliberately deviating from this route. The reason is not difficult to appreciate; at such high values of  $\mu$  the first term in the loading form (62) is independent of the wrinkle mode number  $m$  according to the predictions of stage II. Thus, a simple one-term form for  $\lambda$  would no longer be adequate to capture any wavenumber variation whatsoever, which forces our consideration of the more complicated (62). Now, across the main part of the plate, the series (51) becomes

$$\Phi = \Delta^4 \Phi_0 + \Delta^3 \Phi_1 + \Delta^2 \Phi_2 + \dots \quad \text{and} \quad \Theta = \Delta^2 \Theta_0 + \Delta \Theta_1 + \Theta_2 + \dots,$$

519 where  $\Phi_j \equiv \Phi_j(X)$  and  $\Theta_j \equiv \Theta_j(X)$  ( $j = 0, 1, 2, \dots$ ) are to be determined. Note that  
 520 if we write  $\Phi_0 = \tilde{\mu}^2 \phi_0$ , then  $\phi_0$  satisfies the equation (53) with the value  $\Gamma = K_0$ , by  
 521 virtue of which we are guaranteed that  $\Phi_0'(1) = 0$ . Thus, we are able to express the  
 522 form of the base state in the  $X = \mathcal{O}(1)$  rim region to obtain the counterpart to (54)

523 in the form

$$524 \quad (63a) \quad \Phi = \tilde{\mu}^2 \Delta^4 + (A_{22} X^2 + A_{21} X) \Delta^2 + (A_{13} X^3 + A_{12} X^2 + A_{11} X) \Delta + \dots,$$

$$525 \quad (63b) \quad \Theta = B_{20} \Delta^2 + (B_{11} X + B_{10}) \Delta + \dots,$$

527 where

$$528 \quad A_{22} := \frac{1}{2} \left( 1 - \frac{1}{2} K_0^2 \right) \tilde{\mu}^2, \quad A_{13} := \frac{1}{6} (3 + K_0^2) \tilde{\mu}^2, \quad B_{20} := -K_0 \tilde{\mu}^3,$$

$$529 \quad A_{21} := \frac{\tilde{\lambda}_1 K_1}{\tilde{\mu}}, \quad A_{12} := -\frac{\tilde{\lambda}_1}{2\tilde{\mu}} (K_0 - K_1), \quad B_{11} := 2K_0 \tilde{\mu}^3,$$

$$530 \quad A_{11} := \frac{\tilde{\lambda}_2 K_1}{\tilde{\mu}} + \frac{\tilde{\lambda}_1^2 K_2}{\tilde{\mu}^4}, \quad B_{10} := -\tilde{\lambda}_1.$$

532 Standard numerical work (which is relegated to the supplementary material) shows  
 533 that  $K_0 \simeq 3.212$ ,  $K_1 \simeq 0.5179$  and  $K_2 \simeq 0.0389$ . In the expression (63a) we note the  
 534 absence of the  $\Delta^3$  term which is a direct consequence of the fact that  $\Phi'_0(1) = 0$ . We  
 535 can use the approximation of the basic state (63) to capture the asymptotic structure  
 536 of the wrinkles. To this end we shall employ the ansatz

$$537 \quad (64) \quad W = \widetilde{W}_0 + \widetilde{W}_1 \Delta^{-1} + \dots \quad \text{and} \quad \Psi = \widetilde{\Psi}_0 \Delta^{-1} + \widetilde{\Psi}_1 \Delta^{-2} + \dots$$

538 The second equation in (8) gives an algebraic constraint,  $\widetilde{M} \widetilde{\Psi}_0 + 2K_0 \tilde{\mu}^3 \widetilde{W}_0 = 0$ ;  
 539 thus, the terms in the expansion (64b) are quite passive and respond to what the  $W_j$   
 540 ( $j = 0, 1, \dots$ ) components need to do. However, use of (64) in (8a) yields

$$541 \quad (65) \quad \frac{d^2 \widetilde{W}_0}{dX^2} - (\alpha X - \beta) \widetilde{W}_0 = 0 \quad \text{with} \quad \alpha := -\frac{2\widetilde{M} A_{22}}{\tilde{\mu}^2}, \quad \beta := -\frac{\widetilde{M}(\widetilde{M} - A_{21})}{\tilde{\mu}^2}.$$

542 We recognise this equation once again as related to an Airy form, and elementary  
 543 algebra shows that a non-trivial solution with  $\widetilde{W} \rightarrow 0$  as  $X \rightarrow 0$  and  $X \rightarrow \infty$  is  
 544 possible if

$$545 \quad (66) \quad \tilde{\lambda}_1 K_1 = \tilde{\mu} \widetilde{M} + \xi_0 \left( \frac{1}{2} K_0^2 - 1 \right)^{2/3} \tilde{\mu}^{5/3} \widetilde{M}^{-1/3},$$

546 where  $\text{Ai}(-\xi_0) = 0$ ,  $\xi_0 \simeq 2.331$ . Now, while the leading-order term in (62) was  
 547 independent of  $\widetilde{M}$ , we observe that  $\tilde{\lambda}_1 \rightarrow \infty$  both as  $\widetilde{M} \rightarrow 0$  and as  $\widetilde{M} \rightarrow \infty$ . Thus,  
 548 we can identify the wavenumber that corresponds to the least loading, and minimizing  
 549  $\tilde{\lambda}_1$  with respect to  $\widetilde{M}$  gives the critical point  $(\widetilde{M}_c, \tilde{\lambda}_{1c}) \simeq (1.6877, 13.0346)$ .

550 We remark that the solution  $\widetilde{W}_0$  does not fulfil all eight of the rim conditions  
 551 (10) prescribed. This merely reflects the fact that the majority of the wrinkle zone is  
 552 governed by a system of order less than eight, which means that not all the constraints  
 553 can be satisfied. This does not present any problem and just points to the fact  
 554 that the  $\mathcal{O}(\Delta^{-1})$  rim zone contains an inner region in which the aforementioned  
 555 requirements can be ensured. The details of this inner zone affect later terms in  
 556 our asymptotic series, in particular, they do influence the form of  $\tilde{\lambda}_2$  in (62). The  
 557 manipulations required are routine but lengthy so, in the interest of brevity, the details

558 of the corresponding analysis are consigned to the supplementary material. Here, we  
 559 simply state the final results

560 (67)  $\lambda_c = 3.212\mu^3 + 13.0346\mu^{5/2} + 54.8417\mu^2 + \dots$       and       $m_c^2 = 1.6877\mu^{3/2} + \dots$

561 The predictions of these last formulae are illustrated in Figure 5, where we compare  
 562 them with some direct numerical simulations of (8)-(10). It is clear that the agreement  
 563 is very good. In particular, in the left window the relative errors range from 10% at  
 564  $\mu = 80$  to 5.7% when  $\mu = 120$  and are merely 2.8% once  $\mu = 200$ . The predictions  
 565 of the critical wavenumber differ from the simulations by about 5% when  $\mu = 180$ ;  
 566 although these relative errors are slightly larger than for the critical loading values  
 567 it should be remembered that the asymptotic result (67b) consists of only one term.  
 568 Better improvement could be expected should further terms in (67b) be developed  
 569 but this simple result is sufficiently accurate that the additional effort necessary to  
 570 extricate higher order terms is arguably not commensurate with the likely marginal  
 improvement in results.




FIG. 5. Comparisons between direct numerical simulations of the boundary-value problem (8)-(10) and the critical values (67) for  $30 \leq \mu \leq 200$ . The markers correspond to the former set of data, while the continuous curves represent the asymptotic results.

571

572 **7. Discussion.** In this article we have endeavoured to provide a detailed  
 573 description of the short-wavelength wrinkle modes that develop in a uniformly stretched  
 574 weakly clamped circular plate subjected to a transverse pressure. Three distinct  
 575 regimes of initial stretching have been identified (see Figure 6); in the first of these  
 576 the eigenmodes are located off the rim of the plate at a location determined by the  
 577 solution of a pair of consistency conditions. As the size of the stretching  $\mu$  increases  
 578 then the wrinkles effectively sit at the rim, where they are governed by the solution of  
 579 a scaled Airy equation. A third regime is suggested in which the leading-order loading  
 580 required for wrinkling loses all dependence on the mode number.




FIG. 6. Schematic of the asymptotic regimes studied. Upper line indicates the size of the correction to the leading order wavenumber  $\mathcal{O}(\Delta^{3/2})$ .

581 At the outset our principal motivation behind this work was to shed light on  
 582 the nature of the asymptotic decoupling of the FvK system found recently in some  
 583 related studies [5, 6]. Although there are a number of non-trivial examples in the liter-  
 584 erature in which the asymptotic decoupling of the FvK nonlinear equations has been  
 585 encountered, for example [10]-[11], it should be emphasised that the nature of this  
 586 phenomenon was actually quite different. Indeed, a close look indicates that the afore-  
 587 mentioned references were concerned with out-of-plane bending perturbations from a  
 588 state of plane stress. As a consequence, the compatibility relation in the FvK system  
 589 decoupled at leading-order, giving rise to the standard linear bi-harmonic equation  
 590 for the stress function, and this had the effect of turning the equilibrium equation  
 591 into an expression solvable in closed form. So in spite of the fact that the analysis  
 592 was ostensibly nonlinear, those works ended up dealing with a weak nonlinear pertur-  
 593 bation from a linear plane-stress elastic state. By contrast, the situation present in  
 594 our work is exactly the opposite. Here our perturbations take place relatively remote  
 595 from the original flat state of the circular plate; exactly how remote is something  
 596 that is controlled by the nonlinear basic state. This has significant ramifications for  
 597 the subsequent asymptotic analysis as the nature of basic state is one of the critical  
 598 elements in the implementation of our singular perturbation strategies.

599 It is important to appreciate some of the inherent limitations of our results. We  
 600 have been exclusively focused on the onset of wrinkling which is acknowledged as  
 601 being very awkward to observe in the laboratory. While there are numerous valid  
 602 reasons for understanding onset (or near-threshold phenomena [20]), from the practi-  
 603 cal standpoint wrinkles well into the post-buckling regime are much easier to produce.  
 604 In the far-post-buckling situation traditional simplified theories have been developed  
 605 based on tension field theory [21], [22]. The approach taken by tension-field theory is  
 606 in marked contrast to the bifurcation technique adopted here. Tension-field theory in  
 607 some sense smears out the individual wrinkles and seeks to trace the evolution of the  
 608 boundary separating the wrinkled and un-wrinkled areas. As further evidence that the  
 609 post-buckling regime can behave very differently to the onset problem, we note recent  
 610 results that suggest how spatially varying wavenumbers can be dramatically affected  
 611 by increasing the load; see Paulsen *et al.* [23] and Taffetani & Vella [24], to name just

612 two studies of these effects.

613 It is helpful to note that our results need to be considered carefully if general-  
 614 isations to other geometries are contemplated. An obvious question is to ask how  
 615 our work may be applied to annular plates. In our present study the existence of  
 616 the edge instability is contingent upon the presence of compressive stresses near the  
 617 circumference which is guaranteed if the outer edge of the plate is weakly clamped  
 618 or pinned. If there is also uniform stretching applied along the outer circumference  
 619 then one has a handle on the extent of the region of compressive stresses and this is  
 620 the role played by our parameter  $\mu$ . For an annular plate with a traction-free inner  
 621 boundary, weakly clamped along the outer rim, and subjected to uniform stretching  
 622 along that edge there will be no compressive stresses in the annulus according to the  
 623 Lamé solution. If transverse pressure is also applied then the region of compressive  
 624 stresses will be situated near the outer rim and this will be an entirely nonlinear phe-  
 625 nomenon. Haughton & McKay [25] have considered the plane-stress problem for an  
 626 annular membrane in the case of a nonlinear Varga material and with several types  
 627 of boundary conditions. The principal stresses were found to be always tensile if the  
 628 inner boundary is stress free.

629 Our problem here has the feature that the loading intimately ties together the  
 630 basic state with the infinitesimal wrinkle pattern. The usual approach taken in these  
 631 types of problems is to determine the underlying basic state and then adjust the load-  
 632 ing, which plays the role of an eigenvalue, so that non-trivial modes are possible. Here  
 633 the situation is somewhat different. The value of  $\lambda$  plays a pivotal role in the form of  
 634 the basic state so that both this quantity and the perturbation structure really need  
 635 to be developed in tandem. This is the feature that suggested it would be advanta-  
 636 geous to view  $\lambda$  as given and then calculate the associated wrinkle wavenumber. This  
 637 strategy has enabled us to monitor the stability characteristics of the system as the  
 638 in-plane loading varies from completely unstretched right through to a taut geometry.  
 639 Whilst we have been able to implement similar techniques in related situations, we  
 640 believe this is first example where it has proved possible to track the effect of a varying  
 641 physical parameter over such an extended regime. It would be of considerable inter-  
 642 est to know whether the problem we have here is somewhat special in that respect or  
 643 whether the approach has more general applicability.

644 The referees are thanked for their numerous comments which led to improvements  
 645 in the paper.

646

## REFERENCES

- 647 [1] P.G. CIARLET, *Mathematical Elasticity (vol. 2: Theory of Plates)*, North-Holland, Amsterdam,  
 648 1997.
- 649 [2] B. AUDOLY, AND Y. POMEAY, *Elasticity and Geometry: from hair curls to the non-linear*  
 650 *response of shells*, Oxford University Press, Oxford, 2010.
- 651 [3] J.W. HUTCHINSON, AND Z. SUO, *Mixed mode cracking in layered materials*, Adv. Appl. Mech.,  
 652 29 (1991), pp. 63–93.
- 653 [4] C.D. COMAN, AND A.P. BASSOM, *On the nonlinear membrane approximation and edge-*  
 654 *wrinkling*, Int. J. Solids Struct., 82 (2016), pp. 85–94.
- 655 [5] C.D. COMAN, AND A.P. BASSOM, *Singularities and wrinkling: the case of a concentrated force*,  
 656 Int. J. Eng. Sci., 106 (2016), pp. 229–244.
- 657 [6] C.D. COMAN, M.T. MATTHEWS, AND A.P. BASSOM, *Asymptotic phenomena in pressurised*  
 658 *thin films*, Proc. Roy. Soc. Lond. A, 471 (2015), Article number 20150471.
- 659 [7] C.D. COMAN, *Asymmetric bifurcations in a pressurised circular thin plate under initial tension*,  
 660 Mech. Res. Comm., 47 (2013), pp. 11–17.
- 661 [8] C.D. COMAN, AND A.P. BASSOM, *On the wrinkling of a pre-stressed annular thin film in*

- 662            *tension*, J. Mech. Phys. Solids, 55 (2007), pp. 1601–1617.
- 663 [9] C.D. COMAN, AND D.M. HAUGHTON, *Localized wrinkling instabilities in radially stretched*  
664 *annular thin films*, Acta Mech., 55 (2006), pp. 179–200.
- 665 [10] W.H. WITTRICK, *Axisymmetrical bending of a highly stretched annular plate*, Quart. J. Mech.  
666 Appl. Math., 18 (1965), pp. 11–24.
- 667 [11] W.B. FRASER, *Bending of a highly stretched elastic plate containing an eccentrically plate-*  
668 *reinforced circular hole*, Int. J. Solids Structures, 11 (1975), pp. 501–518.
- 669 [12] G. JUNKIN, AND R. DAVIS, *General non-linear plate theory applied to a circular plate with*  
670 *large deflections*. Int. J. Solids Struct., 9 (1972), pp. 84–93.
- 671 [13] J.G. SIMMONDS, AND A. LIBAI, *Asymptotic forms of a simplified version of the non-linear*  
672 *Reissner equations for clamped elastic spherical caps under outward pressure*, Comp.  
673 Mech., 2 (1987) pp. 231–244.
- 674 [14] U. KOMARAGIRI, M.R. BEGLEY, AND J.G. SIMMONDS, *The mechanical response of freestanding*  
675 *circular elastic films under point and pressure loads*, ASME, J. Appl. Mech., 72 (2005),  
676 pp. 203–212.
- 677 [15] V.L. BERDICHEVSKY, *Variational Principles of Continuum Mechanics*, Nauka, Moscow, 1983.
- 678 [16] D.H. HODGES, *Nonlinear Composite Beam Theory*, American Institute of Aeronautics and  
679 Astronautics Inc.. Roston VA, 2006.
- 680 [17] M. TAYLOR, B. DAVIDOVITCH, Z. QIU, AND K. BERTOLDI, *A comparative analysis of numerical*  
681 *approaches to the mechanics of elastic sheets*, J. Mech. Phys. Solids, 79 (2015), pp. 92–107.
- 682 [18] G.G. ADAMS, *Elastic wrinkling of a tensioned circular plate using von-Kármán plate theory*,  
683 ASME J. Appl. Mech., 60 (1993), pp. 520–525.
- 684 [19] H. TROGER, AND A. STEINDL, *Nonlinear Stability and Bifurcation Theory*, Springer-Verlag,  
685 Wien, 1991.
- 686 [20] B. DAVIDOVITCH, R.D. SCHROLL, D. VELLA, M. ADDA-BEDIA, AND E.A. CERDA, *Prototypical*  
687 *model for tensional wrinkling in thin sheets*, Proc. National Academy Sciences, 108 (2011),  
688 pp. 18227–18232.
- 689 [21] A.C. PIPKIN, *The relaxed energy density for isotropic elastic membranes*, IMA J. appl. Math.,  
690 36 (1986), pp. 85–99.
- 691 [22] D.J. STEIGMANN, *Tension-Field theory*, Proc. R. Soc. Lond. A 429 (1990), pp. 141–173.
- 692 [23] J.D. PAULSEN, E. HOHLFELD, H. KING, J. HUANG, Z. QIU, T.P. RUSSELL, N. MENON, D.  
693 VELLA, AND B. DAVIDOVITCH, *Curvature-induced stiffness and the spatial variation of*  
694 *wavelength in wrinkled sheets*, Proc. National Academy Sciences, 113 (2016), pp. 1144–  
695 1149.
- 696 [24] M. TAFFETANI, AND D. VELLA, *Regimes of wrinkling in pressurized elastic shells*, Phil. Trans  
697 Roy. Soc. Lond. A, 375 (2017). Article number 20160330.
- 698 [25] D.M. HAUGHTON, AND A.B. MCKAY, *Wrinkling of annular discs subjected to radial displace-*  
699 *ments*, Int. J. Engng. Sci., 33 (1995), pp. 335–350.

**SUPPLEMENTARY MATERIALS: WRINKLING STRUCTURES AT  
THE RIM OF AN INITIALLY STRETCHED CIRCULAR THIN  
PLATE SUBJECTED TO TRANSVERSE PRESSURE\***

CIPRIAN D. COMAN<sup>†</sup> AND ANDREW P. BASSOM<sup>‡</sup>

**SM1. Background.** For easy reference we start by listing below the main equations from [SM1]. The basic state is described by the main fields  $\Theta \equiv \Theta(\rho)$  and  $\Phi \equiv \Phi(\rho)$ , which satisfy the nonlinear equations

$$(SM1) \quad \frac{d^2\Theta}{d\rho^2} + \frac{1}{\rho} \frac{d\Theta}{d\rho} - \frac{\Theta}{\rho^2} = \lambda\rho + \frac{\Theta\Phi}{\rho} \quad \text{and} \quad \frac{d^2\Phi}{d\rho^2} + \frac{1}{\rho} \frac{d\Phi}{d\rho} - \frac{\Phi}{\rho^2} = -\frac{\Theta^2}{2\rho},$$

subject to the constraints

$$(SM2a) \quad \Theta(1) = 0, \quad \Phi(1) = \mu^2,$$

$$(SM2b) \quad \Theta(0) = 0, \quad \Phi(0) = 0.$$

The incremental radial amplitudes  $(W, \Psi)$  satisfy two coupled linear bifurcation equations,

$$(SM3) \quad \mathcal{L}_{11}[W] + \mathcal{L}_{12}[\Psi] = 0 \quad \text{and} \quad \mathcal{L}_{21}[W] + \mathcal{L}_{22}[\Psi] = 0,$$

where

$$(SM4a) \quad \mathcal{L}_{11} \equiv [\mathcal{L}_0^{(m)}]^2 - \frac{1}{\rho} \frac{d}{d\rho} \left( \Phi \frac{d}{d\rho} \right) + \frac{d\Phi}{d\rho} \left( \frac{m}{\rho} \right)^2, \quad \mathcal{L}_{22} \equiv [\mathcal{L}_0^{(m)}]^2,$$

$$(SM4b)$$

$$\mathcal{L}_{12} = -\mathcal{L}_{21} \equiv -\frac{1}{\rho} \frac{d}{d\rho} \left( \Theta \frac{d}{d\rho} \right) + \frac{d\Theta}{d\rho} \left( \frac{m}{\rho} \right)^2, \quad \mathcal{L}_0^{(k)} \equiv \frac{1}{\rho} \frac{d}{d\rho} \left( \rho \frac{d}{d\rho} \right) - \frac{k^2}{\rho^2}.$$

The corresponding boundary conditions correspond to a weakly clamped plate and assume the form

$$(SM5) \quad W = \frac{dW}{d\rho} = \Psi = \frac{d\Psi}{d\rho} = 0, \quad \text{for } \rho \in \{0, 1\}.$$

**SM2. Basic state.** Let us recall the main scalings from §6 in [SM1],

$$(SM6) \quad \mu = \Delta^2 \tilde{\mu}, \quad \tilde{\mu} = \mathcal{O}(1),$$

and  $m^2 = \Delta^3 \tilde{M}$ , for some  $\tilde{M} = \mathcal{O}(1)$ ; also, our loading can be expressed as

$$(SM7) \quad \lambda = K_0 \tilde{\mu}^3 \Delta^6 + \tilde{\lambda}_1 \Delta^5 + \tilde{\lambda}_2 \Delta^4 + \dots,$$

for some  $\tilde{\lambda}_j = \mathcal{O}(1)$  ( $j = 1, 2, \dots$ ). Our main goal is to find  $\tilde{\lambda}_1$  and  $\tilde{\lambda}_2$ .

---

\*Submitted to the editors January 18, 2018.

<sup>†</sup>School of Mathematical Sciences, University of Nottingham, University Park, Nottingham NG7 2RD, UK ([cdc3p@yahoo.com](mailto:cdc3p@yahoo.com))

<sup>‡</sup>School of Physical Sciences, University of Tasmania, Private Bag 37, Hobart TAS 7001, Australia ([andrew.bassom@utas.edu.au](mailto:andrew.bassom@utas.edu.au))



A little more than a simple exercise in elementary algebra indicates that away from the rim of the plate ( $\rho = 1$ ) our basic-state fields must be expanded according to

$$(SM8a) \quad \Phi = \Delta^4 \Phi_0 + \Delta^3 \Phi_1 + \Delta^2 \Phi_2 + \dots,$$

$$(SM8b) \quad \Theta = \Delta^2 \Theta_0 + \Delta \Theta_1 + \Theta_2 + \dots,$$

where the behaviours of the unknown coefficient functions  $\Theta_j \equiv \Theta_j(\rho)$  and  $\Phi_j \equiv \Phi_j(\rho)$  ( $j = 0, 1, 2, \dots$ ) can be found as explained below.

Substituting (SM8) in (SM1a) leads to the algebraic relations

$$(SM9a) \quad -K_0 \tilde{\mu}^3 \rho^2 = \Theta_0 \Phi_0,$$

$$(SM9b) \quad -\tilde{\lambda}_1 \rho^2 = \Theta_0 \Phi_1 + \Theta_1 \Phi_0,$$

$$(SM9c) \quad -\tilde{\lambda}_2 \rho^2 = \Theta_0 \Phi_2 + \Theta_1 \Phi_1 + \Theta_2 \Phi_0,$$

while the other base equation, (SM1b), yields a sequence of differential equations

$$(SM10a) \quad \mathcal{L}_0^{(1)}[\Phi_0] = -\frac{\Theta_0^2}{2\rho},$$

$$(SM10b) \quad \mathcal{L}_0^{(1)}[\Phi_1] = -\frac{\Theta_0 \Theta_1}{\rho},$$

$$(SM10c) \quad \mathcal{L}_0^{(1)}[\Phi_2] = -\frac{\Theta_1^2 + 2\Theta_0 \Theta_2}{2\rho}.$$

**SM2.1. Zeroth order.** Eliminating  $\Theta_0$  between (SM9a) and (SM10a), and further putting  $\Phi_0 =: \tilde{\mu}^2 \phi_0$ , gives that

$$(SM11) \quad \phi_0'' + \frac{1}{\rho} \phi_0' - \frac{1}{\rho^2} \phi_0 = -\frac{K_0^2 \rho^3}{2\phi_0^2}, \quad \phi_0(0) = 0, \quad \phi_0(1) = 1, \quad \phi_0'(1) = 0.$$

We recall that the base condition  $\Phi = \mu^2$  on  $\rho = 1$  leads to the middle of the above boundary conditions, while the vanishing of the derivative at the rim was provoked by the outcome of stage II. By solving numerically the eigenvalue problem (SM11) we find

$$(SM12) \quad K_0 \simeq 3.212.$$

We are going to need the form of the basic state inside the rim zone (i.e., the wrinkling layer), where  $\rho = 1 - (X/\Delta)$  with  $X = \mathcal{O}(1)$ . To this end we need to note that  $\phi_0'(1) = 0$  (fixed),  $\phi_0''(1) = 1 - (K_0^2/2)$  (using the equation) and  $\phi_0'''(1) = -(3 + K_0^2)$  (differentiating the equation). Put together these results tell us that, where  $X = \mathcal{O}(1)$ , we have

$$(SM13) \quad \Phi_0 = \tilde{\mu}^2 \left[ 1 + \frac{1}{4}(2 - K_0^2) \left( \frac{X}{\Delta} \right)^2 + \frac{1}{6}(3 + K_0^2) \left( \frac{X}{\Delta} \right)^3 + \dots \right].$$

**SM2.2. First order.** At the next order, eliminating the  $\Theta_0$  and  $\Theta_1$  from the base equations (SM9b) and (SM10b) gives the equation for  $\Phi_1$ . By setting  $\Phi_1 =: \tilde{\mu}^2 \phi_1$  we obtain

$$(SM14) \quad \phi_1'' + \frac{1}{\rho} \phi_1' - \frac{1}{\rho^2} \phi_1 - \frac{K_0^2 \rho^3}{\phi_0^3} \phi_1 = -\frac{K_0 \tilde{\lambda}_1 \rho^3}{\tilde{\mu}^3 \phi_0^2}, \quad \phi_1(0) = 0, \quad \phi_1(1) = 0.$$

Again, we need the Taylor series of  $\phi_1$  where  $X = \mathcal{O}(1)$ . If we define the auxiliary problem

$$(SM15) \quad \tilde{\phi}_1'' + \frac{1}{\rho} \tilde{\phi}_1' - \frac{1}{\rho^2} \tilde{\phi}_1 - \frac{K_0^2 \rho^3}{\phi_0^3} \tilde{\phi}_1 = \frac{K_0 \rho^3}{\phi_0^2}, \quad \tilde{\phi}_1(0) = 0, \quad \tilde{\phi}_1(1) = 0,$$

then this is well-defined and with no parameters, as  $K_0$  is already known. Standard numerical methods help us to identify  $\tilde{\phi}_1'(1) =: K_1$ , namely,

$$(SM16) \quad K_1 \simeq 0.5179.$$

Setting  $\rho \rightarrow 1$  in the differential equation (SM15) also tells us that  $\tilde{\phi}_1''(1) = K_0 - K_1$ . Thus, the Taylor expansion of  $\tilde{\phi}_1$  as  $\rho \rightarrow 1$  can be expressed in the form

$$\tilde{\phi}_1 \rightarrow K_1 \left( -\frac{X}{\Delta} \right) + \frac{1}{2} (K_0 - K_1) \left( \frac{X}{\Delta} \right)^2 + \dots,$$

so that

$$(SM17) \quad \Phi_1 = -\frac{\tilde{\lambda}_1}{\tilde{\mu}} \left[ K_1 \left( -\frac{X}{\Delta} \right) + \frac{1}{2} (K_0 - K_1) \left( \frac{X}{\Delta} \right)^2 + \dots \right].$$

**SM2.3. Second order.** If we repeat the above procedures on (SM9c) and (SM10c) we find that, if  $\Phi_2 =: \tilde{\mu}^2 \phi_2$ , then  $\phi_2$  satisfies

$$\phi_2'' + \frac{1}{\rho} \phi_2' - \frac{1}{\rho^2} \phi_2 - \frac{K_0^2 \rho^3}{\phi_0^3} \phi_2 = -\frac{K_0 \rho^3 \tilde{\lambda}_2}{\tilde{\mu}^3 \phi_0^2} - \frac{\tilde{\lambda}_1^2 \rho^3}{2 \tilde{\mu}^6 \phi_0^2} - \frac{3 K_0^2 \rho^3 \tilde{\lambda}_1^2 \tilde{\phi}_1^2}{2 \tilde{\mu}^6 \phi_0^4} - \frac{2 \tilde{\lambda}_1^2 K_0 \rho^3 \tilde{\phi}_1}{\tilde{\mu}^6 \phi_0^3},$$

and restoring the original variables gives

$$(SM18) \quad \Phi_2'' + \frac{1}{\rho} \Phi_2' - \frac{1}{\rho^2} \Phi_2 - \frac{K_0^2 \rho^3}{\phi_0^3} \Phi_2 = -\frac{\tilde{\lambda}_2}{\tilde{\mu}} \left( \frac{\rho^3 K_0}{\phi_0^2} \right) + \frac{\tilde{\lambda}_1^2}{\tilde{\mu}^4} \left[ -\frac{\rho^3}{2 \phi_0^4} (\phi_0 + K_0 \tilde{\phi}_1) (\phi_0 + 3 K_0 \tilde{\phi}_1) \right].$$

This must be solved subject to the homogeneous boundary conditions  $\Phi_2(0) = \Phi_2(1) = 0$ .

We can take advantage of the linearity of (SM18) and use the principle of superposition to solve it. The particular form of its right-hand side suggests introducing the auxiliary problems

$$(SM19) \quad \Phi_{2a}'' + \frac{1}{\rho} \Phi_{2a}' - \frac{1}{\rho^2} \Phi_{2a} - \frac{K_0^2 \rho^3}{\phi_0^3} \Phi_{2a} = \frac{\rho^3 K_0}{\phi_0^2}, \quad \Phi_{2a}(0) = \Phi_{2a}(1) = 0,$$

and

$$(SM20) \quad \Phi_{2b}'' + \frac{1}{\rho} \Phi_{2b}' - \frac{1}{\rho^2} \Phi_{2b} - \frac{K_0^2 \rho^3}{\phi_0^3} \Phi_{2b} = \frac{\rho^3}{2 \phi_0^4} (\phi_0 + K_0 \tilde{\phi}_1) (\phi_0 + 3 K_0 \tilde{\phi}_1), \quad \Phi_{2b}(0) = \Phi_{2b}(1) = 0.$$

Note that  $\Phi_2 = \Phi_{2a} + \Phi_{2b}$  and we have already dealt with (SM19) in §SM2.2. Direct numerical integration of (SM20) immediately allows us to find  $\Phi'_{2b}(1) =: K_2$ , namely,

$$(SM21) \quad K_2 \simeq 0.0389.$$

Putting this together suggests that as  $\rho \rightarrow 1$  so

$$(SM22) \quad \Phi_2 = \left( K_1 \frac{\tilde{\lambda}_2}{\tilde{\mu}} + K_2 \frac{\tilde{\lambda}_1^2}{\tilde{\mu}^4} \right) \left( \frac{X}{\Delta} \right) + \dots$$

If we then combine the results (SM13), (SM17) and (SM22), we conclude that, where  $X = \mathcal{O}(1)$ , the base-state variable  $\Phi$  assumes the behaviour

$$(SM23) \quad \Phi = \Delta^4 \tilde{\mu}^2 + \Delta^2 \left[ \frac{1}{4} (2 - K_0^2) \tilde{\mu}^2 X^2 + \frac{\tilde{\lambda}_1 K_1}{\tilde{\mu}} X \right] \\ + \Delta \left[ \frac{1}{6} (3 + K_0^2) \tilde{\mu}^2 X^3 + \frac{\tilde{\lambda}_1 (K_1 - K_0)}{2\tilde{\mu}} X^2 + \left( K_1 \frac{\tilde{\lambda}_2}{\tilde{\mu}} + K_2 \frac{\tilde{\lambda}_1^2}{\tilde{\mu}^4} \right) X \right] + \mathcal{O}(1).$$

For simplicity, we shall define a sequence of constants  $A_{ij}$  ( $i, j \in \{1, 2, 3\}$ ) so that this behaviour can be expressed more succinctly as

$$(SM24) \quad \Phi = \tilde{\mu}^2 \Delta^4 + [A_{22} X^2 + A_{21} X] \Delta^2 + [A_{13} X^3 + A_{12} X^2 + A_{11} X] \Delta + \mathcal{O}(1).$$

**SM2.4. The bifurcation equation.** We can use the information contained in (SM24) to deduce that for a wrinkle structure

$$(SM25) \quad W = \tilde{W}_0 + \Delta^{-1} \tilde{W}_1 + \dots \quad \text{and} \quad \Psi = \Delta^{-1} \tilde{\Psi}_0 + \Delta^{-2} \tilde{\Psi}_1 + \dots,$$

for some  $W_j, \Psi_j$  (unknown at this stage). We note that at leading order in (SM3b) we just get

$$\tilde{M} \tilde{\Psi}_0 + 2K_0 \tilde{\mu}^3 \tilde{W}_0 = 0;$$

thus, the  $\Psi_j$ -functions are quite passive and respond to what the  $W_j$ -components need to do. At zeroth orders in the other equation ( $\mathcal{O}(\Delta^6)$ ) we find that

$$(SM26) \quad \mathcal{L}_\#[\tilde{W}_0] \equiv \frac{d^2 \tilde{W}_0}{dX^2} - [\alpha X - \beta] \tilde{W}_0 = 0,$$

where

$$(SM27) \quad \alpha := -\frac{2A_{22} \tilde{M}}{\tilde{\mu}^2} \quad \text{and} \quad \beta := \frac{(A_{21} - \tilde{M}) \tilde{M}}{\tilde{\mu}^2}.$$

The solution of this equation is

$$\tilde{W}_0 \propto \text{Ai}(\alpha^{1/3}(X - \beta/\alpha)),$$

which vanishes as  $X \rightarrow 0$  if  $-\beta/\alpha^{2/3} = -\zeta_0$ , where  $\zeta_0 \simeq 2.331$ . Making use of (SM27) this simplifies to

$$(SM28) \quad \tilde{\lambda}_1 K_1 = \tilde{\mu} \tilde{M} + 2.331 \left( \frac{1}{2} K_0^2 - 1 \right)^{2/3} \tilde{\mu}^{5/3} \tilde{M}^{-1/3},$$

which tells us that  $\tilde{\lambda}_1 = \tilde{\lambda}_1(\tilde{M})$  has the property that  $\tilde{\lambda}_1 \rightarrow +\infty$  as either  $\tilde{M} \rightarrow \infty$  or  $\tilde{M} \rightarrow 0^+$ . Clearly, this indicates that the curve  $\tilde{\lambda}_1$  vs.  $\tilde{M}$  has a minimum,  $(\tilde{M}_c, \tilde{\lambda}_{1c})$  (say), and simple numerical calculations yield

$$(SM29) \quad \tilde{M}_c \simeq 1.6877 \quad \text{and} \quad \tilde{\lambda}_{1c} \simeq 13.0346.$$

**SM2.5. The higher-order correction term.** At  $\mathcal{O}(\Delta^5)$  in (SM3a) we obtain

$$(SM30) \quad -X\tilde{\mu}^2 \frac{d^2\tilde{W}_0}{dX^2} - \tilde{\mu}^2 \frac{d^2\tilde{W}_1}{dX^2} - 2\tilde{M} \frac{d^2\tilde{W}_0}{dX^2} + 2\tilde{M}X(\tilde{M} - 2A_{22}X - A_{21})\tilde{W}_0 \\ + \tilde{M}(2\tilde{M}X - 3A_{13}X^2 - 2A_{12}X - A_{11})\tilde{W}_0 + \tilde{M}(\tilde{M} - 2A_{22}X - A_{21})\tilde{W}_1 = 0.$$

If we recall the definitions of  $\alpha$  and  $\beta$  from (SM27) and work on the right-hand side of the above equation by using the governing equation for  $\tilde{W}_0$ , we can re-cast (SM30) in the simplified form

$$(SM31) \quad \mathcal{L}_\#[\tilde{W}_1] = X \frac{d^2\tilde{W}_0}{dX^2} - \frac{2\tilde{M}}{\tilde{\mu}^2} \frac{d^2\tilde{W}_0}{dX^2} + \frac{\tilde{M}}{\tilde{\mu}^2} (2\tilde{M}X - 3A_{13}X^2 - 2A_{12}X - A_{11})\tilde{W}_0.$$

We only really need to work out how  $\tilde{W}_1$  behaves as  $X \rightarrow 0$ . However, before we can do that it is necessary to simplify further the right-hand side of (SM31). To this end, let us start by noting that equation (SM26) tells us that  $X\tilde{W}_0 = (\tilde{W}_0'' + \beta\tilde{W}_0)/\alpha$ . Thus, reducing the  $X^2\tilde{W}_0$  by replacing one  $X\tilde{W}_0$  in this way gives

$$(SM32) \quad \mathcal{L}_\#[\tilde{W}_1] = \left(1 - \frac{3\tilde{M}A_{13}}{\tilde{\mu}^2\alpha}\right) X \frac{d^2\tilde{W}_0}{dX^2} - \frac{2\tilde{M}}{\tilde{\mu}^2} \frac{d^2\tilde{W}_0}{dX^2} \\ + \left[\frac{2\tilde{M}(\tilde{M} - A_{12})}{\tilde{\mu}^2} - \frac{3\beta\tilde{M}A_{13}}{\tilde{\mu}^2\alpha}\right] X\tilde{W}_0 - \frac{\tilde{M}A_{11}}{\tilde{\mu}^2}\tilde{W}_0.$$

By differentiating  $X\tilde{W}_0 = (\tilde{W}_0'' + \beta\tilde{W}_0)/\alpha$  we have that  $X\tilde{W}_0' = ((\tilde{W}_0''' + \beta\tilde{W}_0')/\alpha) - \tilde{W}_0$  and  $X\tilde{W}_0'' = ((\tilde{W}_0'''' + \beta\tilde{W}_0'')/\alpha) - 2\tilde{W}_0'$ . Thus, the right-hand side of (SM32) becomes

$$(SM33) \quad RHS := \left(1 - \frac{3\tilde{M}A_{13}}{\tilde{\mu}^2\alpha}\right) \left[\frac{1}{\alpha}(\tilde{W}_0'''' + \beta\tilde{W}_0'') - 2\tilde{W}_0'\right] - \frac{2\tilde{M}}{\tilde{\mu}^2} \frac{d^2\tilde{W}_0}{dX^2} \\ + \left[\frac{2\tilde{M}(\tilde{M} - A_{12})}{\tilde{\mu}^2} - \frac{3\beta\tilde{M}A_{13}}{\tilde{\mu}^2\alpha}\right] \frac{1}{\alpha}(\tilde{W}_0'' + \beta\tilde{W}_0) - \frac{\tilde{M}A_{11}}{\tilde{\mu}^2}\tilde{W}_0.$$

Now the *RHS* is expressed as a linear multiple of various derivatives of  $\tilde{W}_0$ . To write down the solution of  $\mathcal{L}_\#[\tilde{W}_1] = RHS$ , we need the following observation. If we denote by  $g^{(n)}$  the  $n^{\text{th}}$ -order derivative of the function  $g \equiv g(X)$  ( $n \in \mathbb{N}$ ), then the particular integral of the equation in  $f \equiv f(X)$ ,

$$\mathcal{L}_\#[f] = g^{(n)},$$

is given by

$$f = f_{\text{part}}(X) := \frac{1}{(n+1)\alpha} g^{(n+1)}(X).$$

Given these observations we can now write down the solution of the full  $\widetilde{W}_1$  equation. Putting everything together, we finally get

$$(SM34) \quad \widetilde{W}_1 = \left(1 - \frac{3\widetilde{M}A_{13}}{\widetilde{\mu}^2\alpha}\right) \left[\frac{1}{\alpha^2} \left(\frac{1}{5}\widetilde{W}_0^{(5)} + \frac{1}{3}\beta\widetilde{W}_0''''\right) - \frac{1}{\alpha}\widetilde{W}_0''\right] - \frac{2\widetilde{M}}{\widetilde{\mu}^2} \frac{\widetilde{W}_0'''}{3\alpha} \\ + \left[\frac{2\widetilde{M}(\widetilde{M} - A_{12})}{\widetilde{\mu}^2} - \frac{3\beta\widetilde{M}A_{13}}{\widetilde{\mu}^2\alpha}\right] \frac{1}{\alpha^2} \left(\frac{1}{3}\widetilde{W}_0'''' + \beta\widetilde{W}_0'\right) - \frac{\widetilde{M}A_{11}}{\widetilde{\mu}^2\alpha} \widetilde{W}_0'.$$

Let us recall that we are solely interested in what happens to (SM34) as  $X \rightarrow 0$ . We already know that  $\widetilde{W}_0(0) = 0$ , and we set  $\widetilde{W}_0'(0) =: \omega_0$ . In light of this notation the governing equation (SM26) and its differential consequences imply that

$$\widetilde{W}_0''(0) = 0, \quad \widetilde{W}_0'''(0) = -\beta\omega_0, \quad \widetilde{W}_0^{(4)}(0) = 2\alpha\omega_0, \quad \widetilde{W}_0^{(5)}(0) = \beta^2\omega_0.$$

Together with (SM34) this then leads us to (SM35)

$$\widetilde{W}_1 \rightarrow \left[-\frac{2\beta^2}{15\alpha^2} - \frac{8\beta^2 A_{13}\widetilde{M}}{5\widetilde{\mu}^2\alpha^3} + \frac{2\widetilde{M}\beta}{3\alpha\widetilde{\mu}^2} + \frac{4\widetilde{M}\beta(\widetilde{M} - A_{12})}{3\alpha^2\widetilde{\mu}^2} - \frac{\widetilde{M}A_{11}}{\alpha\widetilde{\mu}^2}\right] \omega_0, \quad \text{as } X \rightarrow 0.$$

**SM3. The bending layer.** To tie things down we still need to consider the rim bending layer where all the boundary conditions on the perturbation are imposed. It can be shown by easy balances that the depth of the inner zone is  $\mathcal{O}(\Delta^{-2})$ , so we are led to introduce a new rescaled variable  $\zeta = \mathcal{O}(1)$  defined by

$$\rho = 1 - \frac{\zeta}{\Delta^2}.$$

We are somewhat fortunate as this happens to be the rim layer for the base state as well. This layer only operates on the  $\Theta$  component and that is just too small to come into play (so the driving differential operator in the bending layer will have constant coefficients). It turns out that the leading-order equation for the  $W$ -component of the wrinkle is just

$$\frac{d^4 W_{\text{bend}}}{d\zeta^4} - \mu^2 \frac{d^2 W_{\text{bend}}}{d\zeta^2} = 0.$$

We need the solution of this differential equation to match onto the linearly decaying  $\widetilde{W}_0$  as  $\zeta \rightarrow \infty$ , and to satisfy the rim conditions that  $W_{\text{bend}}$  and its first derivative vanish on  $\zeta = 0$ ; these constraints leave us with

$$W_{\text{bend}} = \zeta + \frac{1}{\widetilde{\mu}} \exp(-\widetilde{\mu}\zeta) - \frac{1}{\widetilde{\mu}}.$$

So this tells us that  $W_{\text{bend}}$  grows like  $\zeta$ , while the constant part of its large- $\zeta$  behaviour is simply  $-1/\widetilde{\mu}$ . We can now take advantage of these observations in conjunction with (SM35) to deduce that

$$(SM36) \quad -\frac{2\beta^2}{15\alpha^2} - \frac{8\beta^2 A_{13}\widetilde{M}}{5\widetilde{\mu}^2\alpha^3} + \frac{2\widetilde{M}\beta}{3\alpha\widetilde{\mu}^2} + \frac{4\widetilde{M}\beta(\widetilde{M} - A_{12})}{3\alpha^2\widetilde{\mu}^2} - \frac{\widetilde{M}A_{11}}{\alpha\widetilde{\mu}^2} = -\frac{1}{\widetilde{\mu}},$$

whence, by re-arrangement,

$$(SM37) \quad \widetilde{\lambda}_2 = \frac{\widetilde{\mu}}{K_1} \left[ \frac{1}{\widetilde{M}} \left( \widetilde{\mu}\alpha - \frac{2\beta^2\widetilde{\mu}^2}{15\alpha} \right) - \frac{8A_{13}\beta^2}{5\alpha^2} + \frac{2\beta}{3} + \frac{4\beta(\widetilde{M} - A_{12})}{3\alpha} - \frac{\widetilde{\lambda}_1^2 K_2}{\widetilde{\mu}^4} \right].$$

Since by our original assumption (SM6)  $\tilde{\mu} = \mathcal{O}(1)$ , we are free to set  $\tilde{\mu} = 1$  in (SM37). Substituting also the numerical values (SM12), (SM16), (SM21) and (SM29) we eventually get

$$(SM38) \quad \tilde{\lambda}_2 \simeq 54.8417.$$

## REFERENCES

- [SM1] C.D. COMAN, AND A.P. BASSOM, *Wrinkling structures at the rim of an initially stretched circular thin plate subjected to transverse pressure*, Submitted.