

Reference

- Aerts S., Van Loo, P., Thijs G., Moreau Y. and De Moor B.** (2003) Computational detection of cis-regulatory modules. *Bioinformatics* 19: II5-II14.
- Aharoni A., Dixit S., Jetter R., Thoenes E., Arkel G. V. and Pereira A.** (2004) The SHINE Clade of AP2 Domain Transcription Factors Activates Wax Biosynthesis, Alters Cuticle Properties, and Confers Drought Tolerance when Overexpressed in *Arabidopsis*. *The Plant Cell* 16:2463-2480.
- Alonso JM. And Stepanova AN.** (2004) The Ethylene Signaling Pathway. *Science* 306(5701): 1513-1515.
- Andeme-Onzighi C., Sivaguru M., Judy-March J., Baskin T. I. and Driouich A. (2002)** The *reb1-1* mutation of *Arabidopsis* alters the morphology of trichoblasts, the expression of arabinogalactan-proteins and the organization of cortical microtubules. *Planta* 215: 949-958.
- Baluska F., Salaj J., Mathur J., Braun M., Jasper F., Amaj J., Chua N., Barlow P. W. and Volkmann D.** (2000) Root Hair Formation: F-Actin-Dependent Tip Growth Is Initiated by Local Assembly of Profilin-Supported F-Actin Meshworks Accumulated within Expansin-Enriched Bulges. *Developmental Biology* 227(2): 618-632.
- Bannister J.V., Bannister W.H., Rotilio G.** (1987) Aspects of the structure, function, and applications of superoxide dismutase. *CRC Crit Rev Biochem* 22: 111-180.
- Bates T. R. and Lynch J. P.** (2000) Plant growth and phosphorus accumulation of wild type and two root hair mutants of *Arabidopsis thaliana* (Brassicaceae). *American Journal of Botany* 87:958-963.

Belfield E. J., Ruperti B., Roberts J. A. and McQueen-Mason S. (2005) Changes in expansin activity and gene expression during ethylene-promoted leaflet abscission in *Sambucus nigra*. *Journal of Experiment Botany* 56: 817-823.

Belfield, E.J., Ruperti, B., Roberts, J.A. and McQueen-mason, S. (2005) Changes in expansin activity and gene expression during ethylene-promoted leaflet abscission in *Sambucus nigra*. *Journal of Experimental Botany* 56 (413): 817-823.

Bibikova T. N., Blancaflor E. B. and Gilroy S. (1999) Microtubules regulate tip growth and orientation in root hairs of *Arabidopsis thaliana*. *Plant Journal* 17(6):657-65.

Bibikova T. N., Jacob T., Dahse I. and Gilroy S. (1998) Localized changes in apoplastic and cytoplasmic pH are associated with root hair development in *Arabidopsis thaliana*. *Development* 125: 2925-2934.

Bisson MM., Bleckmann A., Allekotte S. and Groth G. (2009) EIN2, the central regulator of ethylene signalling, is localised at the ER membrane where it interacts with the ethylene receptor ETR1. *Biochemical Journal* 424: 1-6.

Bleecker A.B., Patterson S.E. (1997). Last exit: senescence, abscission and meristem arrest in *Arabidopsis*. *Plant cell* 9:1169-79.

Braun M., Baluška F., Witsch M. V. and Menzel D. (1999) Redistribution of actin, profilin and phosphatidylinositol-4, 5-bisphosphate in growing and maturing root hairs. *Planta* 209: 435-443.

Bunning E. (1951) Über die Differenzierungsvorgänge in der Cruciferenwurzel. *Planta* 39: 126-53.

Burns JK, Lewandowski DJ, Nairn CJ and Brown GE. (1998) Endo-1,4- β -glucanase gene expression and cell wall hydrolase activities during abscission in Valencia orange. *Physiologia Plantarum* 102:217-225.

Butenko, M. A., Patterson, S.E., Grini, P. E., Stenvik, G. E., Amundsen, S. S., Mandal, A. and Aalen, R. B. (2003). Inflorescence deficient in abscission controls floral organ abscission in Arabidopsis and identifies a novel family of putative ligands in plants. *Plant cell* 15: 2296-2307.

Cai S. and Lashbrook C. C. (2008) Stamen abscission zone transcriptome profiling reveals new candidates for abscission control: enhanced retention of floral organs in transgenic plants overexpressing Arabidopsis ZINC FINGER PROTEIN2. *Plant Physiol.* 146(3):1305-21.

Carol R. J. and Dolan L. (2002) Building a hair: tip growth in *Arabidopsis thaliana* root hairs. *Philosophical Transactions of the Royal Society of London* 357 (1422): 815-821.

Chao Q., Rothenber M., Solano R., Roman G., Terzaghi W., Ecker JR. (1997). Activation of ethylene gas response pathway in Arabidopsis by the nuclear protein ethylene insensitive 3 and related proteins. *Cell* 89:1133-44.

Chen YF, Randlett MD, Findell JL, Schaller GE (2002) Localization of the ethylene receptor ETR1 to the endoplasmic reticulum of Arabidopsis. *J Biol Chem* 277: 19861–19866.

Cho H. T. and Gosgrove D. J. (2000) Altered expression of expansin modulates leaf growth and pedicel abscission in *Arabidopsis thaliana*. *PNS* 97(17): 9783-9788.

Cho SK., Larue CT., Chevalier D., Wang H., Jinn TL., Zhang S. and Walker JC. (2008) Regulation of floral organ abscission in *Arabidopsis thaliana*. *PNS* 105 (40): 15629-15634.

Clough S. J. and Bent A. F. (1998) Floral dip: a simplified method for Agrobacterium-mediated transformation of *Arabidopsis thaliana*. *Plant Journal* 16(6): 735-43

Cormack R. (1935) Investigations on the development of root hairs. *New Phytologist*. 34(1).

Cosgrove, D.J. (2005). Growth of the plant cell wall. *Nature Reviews. Molecular Cell Biology* 6(11): 850-861.

Coupe S. A. and TAYLOR J. (1997) Temporal and spatial expression of mRNAs encoding pathogenesis - related proteins during ethylene - promoted leaflet abscission in *Sambucus nigra*. *Plant, Cell & Environment* 20(12): 1517-1524.

Cutter, E.G. (1978). The epidermis. In *Plant Anatomy* (London: Clowes and Sons) pp. 94-106.

del Campillo E. and Lewis L. N. (1992) Identification and kinetics of accumulation of proteins induced by ethylene in bean abscission zones. *Plant Physiology* 98: 955-961.

Del Degan F., Child R., Svendsen I. and Ulvskov P. (2001) The cleavable N-terminal domain of plant endopolygalacturonases from clade B may be involved in a regulated secretion mechanism. *The Journal of Biological Chemistry* 176: 35297-35304.

Djinovic-Carugo K., Gautel M. Ylänne J. and Young P. (2002) The spectrin repeat: a structural platform for cytoskeletal protein assemblies. *FEBS Lett.* 513(1):119-23. Review.

Dolan L. (1997) The role of ethylene in the development of plant form. *Journal of Experimental Botany* 48: 201-210.

Dolan, L., Duckett, C. M., Grierson, C., Linstead, P., Schneider, K., Lawson, E., Dean, C., Poethig, S. and Roberts, K. (1994). Clonal origin and patterning in the root epidermis of *Arabidopsis*. *Development* 120: 2465-2474.

Durrant W.E. and Dong X. (2004) Systemic acquired resistance. Annual *Review of Phytopathology* 42:185-209.

Ellis C.M., Nagpal P., Young J. C., Hagen G., Guifoyle T. J. and Reed J. W. (2005) AUXIN RESPONSE FACTOR1 and AUXIN RESPONSE FACTOR2 regulate senescence and floral organ abscission in *Arabidopsis thaliana*. *Development* 132: 4563-4574.

Favery B., Ryan E., Foreman J., Linstead P., Boudonck., Steer M., Shaw P. and Dolan L. (2001) KOJAK encodes a cellulose synthase-like protein required for root hair cell morphogenesis in *Arabidopsis*. *Genes & Dev.* 15: 79-89.

Fernandez D. E., Heck G. R., Perry S. E., Patterson S. E., Bleecker A. B., and Fang S. C (2000) The Embryo MADS Domain Factor AGL15 Acts Postembryonically: Inhibition of Perianth Senescence and Abscission via Constitutive Expression. *Plant Cell* 12: 183-198.

Galway M. E., Masucci J. D., Lloyd A. M., Walbot V., Davis R. W. and Schiefelbein J. W. (1994) The *TTG* Gene Is Required to Specify Epidermal Cell Fate and Cell Patterning in the *Arabidopsis* Root. 166(2): 740-754.

Gasteiger E., Gattiker A., Hoogland C., Ivanyi I., Appel R. D. and Bairoch A. (2003) ExPASy: The proteomics server for in-depth protein knowledge and analysis. *Nucleic Acids Res.* 31(13):3784-8.

Gonzalez-Carranza Z. H., Lozoya-Gloria E., Roberts J. A. (1998). Recent developments in abscission: shedding light on the shedding process. *Trends Plant Sci.* 3: 10-14.

Gonzalez-Carranza Z. H., Elliott K. A. and Roberts J. A. (2007). Expression of polygalacturonases and evidence to support their role during cell separation processes in *Arabidopsis thaliana*. *Journal of Experimental Botany* 58: 3719-3730.

Green P. B. (1962) Mechanism for Plant Cellular Morphogenesis. *Science* 138(3548):1404-5.

Grierson C. S., Roberts K., Feldmann K. A. and Dolan L. (1997) The COW1 Locus of *Arabidopsis* Acts after *RHD2*, and in Parallel with *RHD3* and *TIP1*, to Determine the Shape, Rate of Elongation, and Number of Root Hairs Produced from Each Site of Hair Formation. *Plant Physiology* 115 (3): 981-990.

Guo H. and Ecker J. (2004) The ethylene signaling pathway: new insights. *Current Opinion in Plant Biology* 7: 40-49

Ha C. M, Jun J. H., Nam H. G., and Fletcher J. C. (2004) BLADE-ON-PETIOLE1 encodes a BTB/POZ domain protein required for leaf morphogenesis in *Arabidopsis thaliana*, *Plant Cell Physiol* 45 :1361–1370.

Harding E. W., Tang W., Nichols K. W., Fernandez D. E., and Perry S. E. (2003) Expression and Maintenance of Embryogenic Potential Is Enhanced through Constitutive Expression of AGAMOUS-Like 15. *Plant Physiology* 133:653-663.

Harris N., Taylor J. E. and Roberts J. A. (1997) Characterization and expression of an mRNA encoding a wound-induced (Win) protein from ethylene-treated tomato leaf abscission zone tissue. *Journal of Experimental Botany* 48 (6): 1223-1227.

Heath I. B. and Seagull R. W. (1982) Oriented cellulose fibrils and the cytoskeleton: a critical comparison of models. *Academic Press: London* 163-82.

Hepworth, S. R., Zhang, Y. L., Mckim, S., Li, X. and Haughn, G. (2005). BLADE-ON-PETIOLE-dependent signaling controls leaf and floral patterning in *Arabidopsis*. *Plant Cell* 17: 1434-1448.

Hong S. B., Sexton R. and Tucker M. L. (2000) Analysis of gene promoters for two tomato polygalacturonases expressed in abscission zones and the stigma. *Plant Physiology* 123: 869-881.

Honma T. and Goto K. (2001) Complexes of MADS-box proteins are sufficient to convert leaves into floral organs. *Nature* 409: 525-529

Hruz T., Laule O., Szabo G., Wessendorp F., Bleuler S., Oertle L., Widmayer P., Gruissem W. and Zimmermann P. (2008) Genevestigator v3: a reference expression database for the meta-analysis of transcriptomes. *Adv Bioinformatics* 2008:420747.

Hu Y., Zhong R., Morrison W. H. and Ye Z. H. (2003) The *Arabidopsis RHD3* gene is required for cell wall biosynthesis and actin organization. *Planta* 217(6): 912-21.

Hua J., Chang C., Sun Q. and Meyerowitz E.M. (1995) Ethylene insensitivity conferred by *Arabidopsis ERS* gene. *Science* 269: 1712–1714.

Hua J., Sakai S., Nourizadeh S., Chen Q.C., Bleecker A.B., Ecker J.R. and Meyerowitz E.M. (1998) EIN4 and ERS2 are members of the putative ethylene receptor gene family in *Arabidopsis*. *Plant Cell* 10: 1321-1332.

Huber DJ. (1983) The role of cell wall hydrolases in fruit softening. *Hortic. Rev.* 5: 169-215.

Iyer R. P. (2001) Nucleobase protection of deoxyribo- and ribonucleosides. *Curr Protoc Nucleic Acid Chem.* Chapter 2: Unit 2.1.

Jinn, T. L., Stone, J. M. and Walker, J. C. (2000). HAESA, an *Arabidopsis* leucine-rich repeat receptor kinase, controls floral organ abscission. *Genes Dev.* 14: 108-117.

Jones M. A., Raymond M. J. and Smirnoff N. (2005) Analysis of the root-hair morphogenesis transcriptome reveals the molecular identity of six genes with roles in root-hair development in **Arabidopsis**. *The Plant Journal* 45(1): 83–100.

Kalaitzis P, Solomos T and Tucker ML. (1997) Three different polygalacturonases are expressed in tomato leaf and flower abscission, each with a different temporal expression pattern. *Plant Physiology* 113:1303-1308.

Kalaitzis, P., Koehler, S.M. and Tucker, M.L. (1995) Cloning of a tomato polygalacturonase expressed in abscission. *Plant Molecular Biology* 28 (4): 657-656.

Kandasamy M. K., McKinney E. C., Meagher R. B. (2003) Cell cycle-dependent association of *Arabidopsis* actin-related proteins AtARP4 and AtARP7 with the nucleus. *The Plant Journal* 33(5): 939-948.

Kandasamy, M. K., Deal, R. B., McKinney, E. C. and Meagher, R. B. (2005a). Silencing the nuclear actin-related protein AtARP4 in *Arabidopsis* has multiple effects on plant development, including early flowering and delayed floral senescence. *Plant J.* 41: 845-858.

Kandasamy, M. K., Deal, R. B., McKinney, E. C. and Meagher, R. B. (2005b). Arabidopsis ARP7 is an essential actin-related protein required for normal embryogenesis, plant architecture, and floral organ abscission. *Plant Physiol.* 138: 2019-2032.

Karimi, M., De Meyer, B. and Hilson, P. (2005) Modular cloning in plant cells. *Trends Plant Sci.* 10(3): 103-105.

Karin M. (1990) Too many transcription factors: positive and negative interactions. *New Biol.* 2(2):126-31. Review.

Kelley L. A. and Sternberg M. J. (2009) Protein structure prediction on the Web: a case study using the Phyre server. *Nat Protoc* 4(3):363-71.

Kieber J.J., Rothenberg M., Roman G., Feldmann K.A. and Ecker J.A. (1993) *CTR1*, a negative regulator of the ethylene response pathway in *Arabidopsis*, encodes a member of the Raf family of protein kinases. *Cell* 72:427–441.

Kemmerer E. C. and Tucker M. L. (1994) Comparative study of cellulases associated with adventitious root initiation, apical buds, and leaf, flower, and pod abscission zones in soybean. *Plant Physiology* 104(2):557-562.

Krivan W. and Wasserman W.W. (2001) A predictive model for regulatory sequences directing liver-specific transcription. *Genome Res.* 11: 1559–1566.

Kubigsteltig I, Laudert D, Weiler EW. (1999) Structure and regulation of the *Arabidopsis thaliana* allene oxide synthase gene. *Planta* 208: 63-71.

Kumpf R., Shi C., Larrieu A., Butenko M. A., Tandstad N M. and Aalen R. B. (2010) *INFLORESCENCE DEFICIENT IN ABSCISSION (IDA)* is involved in separation of cortex and epidermal cells during lateral root emergence. *21ST INTERNATIONAL CONFERENCE ON ARABIDOPSIS RESEARCH*.

Lanahan M. B., Yen H. C., Giovannoni J. J. and Klee H. J. (1994) The Never Ripe Mutation Blocks Ethylene Perception in Tomato. *THE PLANT CELL* 6:521-530.

Lashbrook C. C., Giovannoni J. J., Hall B. D., Fischer R. L. and Bennett A. B. (1998) Transgenic analysis of tomato endo-beta-1,4-glucanase gene function. Role of *cel1* in floral abscission. *The Plant Journal* 13(3): 303-310.

Lashbrook C. C., Gonzalez-Bosch C. and Bennett A. B. (1994) Two divergent endo-beta-1,4-glucanase genes exhibit overlapping expression in ripening fruit and abscission flowers. *Plant Cell* 6(10): 1485-1493.

Latchman D. S. (1997) Transcription factors: an overview. *Int J Biochem Cell Biol.* 29(12):1305-12. Review.

Lenne P. F., Raae A. J., Altmann S. M., Saraste M. and Hörber J. K. (2000) States and transitions during forced unfolding of a single spectrin repeat. *FEBS Lett.* 476(3):124-8.

Leslie ME., Lewis MW., Youn JY., Daniels MJ. And Liljegren SJ. (2010) The EVERSHED receptor-like kinase modulates floral organ shedding in *Arabidopsis*. *Development* 137: 467-476.

Letunic I., Doerks T. and Bork P. (2009) SMART 6: recent updates and new developments. *Nucleic Acids Res.* 37(Database issue):D229-32.

Levitt M. and Chothia C. (1976) Structural patterns in globular proteins. *Nature* 261(5561):552-8.

Lew P. R. (1991) Electrogenic transport properties of growing *Arabidopsis* root hairs: the plasma membrane proton pump and potassium channels. *Plant physiology* 97(4): 1527-1534.

Lewis L. N. and Varner J. E. (1970) Synthesis of cellulose during abscission of *Phaseolus vulgaris* leaf explants. *Plant Physiology* 46(2): 194-199.

Lewis M.W., Leslie M.E., Liljegren S.J., (2005) Plant separation: 50 ways to leave your mother. *Current Opinion in Plant Biology* 9: 59–65.

Leyser H., Pickett F. B., Dharmasiri S. and Estelle M. (1996) Mutations in the *AXR3* gene of *Arabidopsis* result in altered auxin response including ectopic expression from the SAUR-AC1 promoter. *Plant journal* 10(3): 403-413.

Libertini E., Li Y. and McQueen-Mason S. J. (2004) Phylogenetic analysis of the plant endo-beta-1,4-glucanase gene family. *Journal of Molecular Evolution* 58(5): 506-515.

Liljegren S.J., Ditta G.S., Eshed Y., Savidge B., Bowman J.L., Yanofsky M.F. (2000) SHATTERPROOF MADS-box genes control seed dispersal in *Arabidopsis*. *Nature* 404:766-70.

Liu, Q., Xu, C. and Wen C. K. (2010) Genetic and transformation studies reveal negative regulation of *ERS1* ethylene receptor signalling in *Arabidopsis*. *BMC PLANT BIOLOGY* 10: 60.

Mao L., Begum D, Chuang H., Budiman M.A., Szymkowiak E.J., et al. (2000). JOINTLESS is a MADS-box gene controlling tomato flower abscission zone development. *Nature* 406: 910-13.

Masucci J. D and Schiefelbein J. W. (1994) The *rhd6* mutation of *Arabidopsis thaliana* alters root-hair initiation through an auxin-and ethylene-associated process. *Plant Physiology* 106(4): 1335-1346.

Masucci J. D and Schiefelbein J. W. (1996) Hormones Act Downstream of TTG and GL2 to Promote Root Hair Outgrowth during Epidermis Development in the Arabidopsis Root. *The Plant Cell* 8(9): 1505-1517.

McKim S.M., Stenvik G.E., Buntenko M.A., Kristiansen W., Cho S.K., Hepworth S.R., Aalen R.B. and Haughn G.w. (2007) The BLADE-ON-PETIOLE genes are essential for abscission zone formation in Arabidopsis. *Development* 135: 1537-1546.

Meri S., Hunter D. A., Chen J. C., Halaly V. and Reid M. S. (2006) Molecular changes occurring during acquisition of abscission competence following auxin depletion in *Mirabilis jalapa*. *Plant Physiology* 14: 1604-1616.

Morre D. (1968) Cell wall dissolution and enzyme secretion during leaf abscission. *Plant Physiology* 43: 1545-1559.

Moss A. C., Doran P. P. and MacMathuna P. (2007) *In Silico* Promoter Analysis can Predict Genes of Functional Relevance in Cell Proliferation: Validation in a Colon Cancer Model. *Translational Oncogenomics* 2007: 2 1–16.

Nikolov D. B and Burley S. K. (1997) RNA polymerase II transcription initiation: a structural view. *Proc Natl Acad Sci U S A*. 94(1):15-22.

Ogawa M, Kay P, Wilson S, Swain SM. (2009) ARABIDOPSIS DEHISCENCE ZONE POLYGALACTURONASE1 (ADPG1), ADPG2, and QUARTET2 are Polygalacturonases required for cell separation during reproductive development in Arabidopsis. *Plant Cell*. 21:216-33.

Ogawa M., Kay P., Wilson S. and Swain S. M. (2009) ARABIDOPSIS DEHISCENCE ZONE POLYGALACTURONASE1 (ADPG1), ADPG2, and QUARTET2 are Polygalacturonases required for cell separation during reproductive development in Arabidopsis. *Plant Cell* 21(1):216-33.

Okushima Y., Overvoorde P. J., Arima K., Alonso J. M., Chan A., Chang C., Ecker JR., Hughes B., Lui A., Nguyen D., Onodera C., Quach H., Smith A., Yu G and Theologis A. (2005) Functional genomic analysis of the *AUXIN RESPONSE FACTOR* gene family members in *Arabidopsis thaliana*: unique and overlapping functions of ARF7 and ARF19. *The Plant Cell* 17:444-463.

Palme K. and Gälweiler L. (1999) PIN-pointing the molecular basis of auxin transport. *Current Opinion in Plant Biology* 2(5): 375-381.

Parker J. S., Cavell A. C., Dolan L., Roberts K., and Grierson C. S. (2000) Genetic Interactions during Root Hair Morphogenesis in *Arabidopsis*. *The Plant Cell* 12(10): 1961-1974.

Pascual J., Pfuhl M., Rivas G., Pastore A. and Saraste M. (1996) The spectrin repeat folds into a three-helix bundle in solution. *FEBS Lett.* 383(3):201-7.

Patterson SE (2001) Cutting loose, Abscission and dehiscence in Arabidopsis. *Plant Physiology* 126: 494-500.

Paul. C. M. (1974): Corn: Its Origin, Evolution and Improvement, *Belknap Press*.

Pelaz S., Gustafson-Brown C., Kohalmi SE. Crosby WL. And Yanofsky MF. (2001) *APETALA1* and *SEPALLATA3* interact to promote flower development. *Plant J.* 26(4): 385-394.

Pinyopich, A., Ditta, G.S., Baumann, E., Wisman, E., and Yanofsky, M.F. (2003). Unraveling the redundant roles of MADS-box genes during carpel and fruit development. *Nature* 424: 85–88.

Qiu P., Feng X. H. and Li L. (2003) Interaction of Smad3 and SRF-associated complex mediates TGF-beta1 signals to regulate SM22 transcription during myofibroblast differentiation. *Mol Cell Cardiol.* 35(12):1407-20.

Rahman A., Hosokawa S., Oono Y., Amakawa T., Goto N. and Tsurumi S. (2002) Auxin and Ethylene Response Interactions during Arabidopsis Root Hair Development Dissected by Auxin Influx Modulators. *Plant Physiology* 130(4): 1908-1917.

Rasori A., Ruperti B., Bonghi C., Tonutti P. and Ramina A. (2002) Characterization of two putative ethylene receptor genes expressed during peach fruit development and abscission. *Journal of Experimental Botany* 53 (379): 2333-2339.

Ren B., Robert F., Wyrick J. J., Aparicio O., Jennings E. G., Simon I., Zeitlinger J., Schreiber J., Hannett N., Kanin E., Volkert T. L., Wilson C. J., Bell S. P. and Young R. A. (2000) Genome-wide location and function of DNA binding proteins. *Science* 290(5500):2306-9.

Rick CM. and Butler L. (1956) Cytogenetics of the Tomato. *Advances in Genetics* 8: 267-382.

Rigas S., Debrosses G., Haralampidis K., Vicente-Agullo F., Feldmannd K. A., Grabovc A., Dolan L. Hatzopoulos P. (2001) *TRH1* Encodes a Potassium Transporter Required for Tip Growth in *Arabidopsis* Root Hairs. *Plant Cell* 13: 139-151.

Roberts JA, Elliott KA, Gonzalez-Carranza ZH. (2002) Abscission, dehiscence and other cell separation processes. *Annual Review of Plant Biology* 53: 131-158.

Roberts JA, Whitelaw CA, Gonzalez-Carranza ZH, McManus M. (2000) Cell separation processes in plants – Models. *Annals of Botany* 86: 223-235.

Roeder R. G. (1996) The role of general initiation factors in transcription by RNA polymerase II. *Trends Biochem Sci.* 21(9):327-35. Review.

RYAN E., GRIERSON C., CAVELL A., STEER M. and Dolan L. (1998) *TIP1* is required for both tip growth and non-tip growth in *Arabidopsis*. *New Phytologist* 138(1): 49-58.

Sakai H., Hua J., Chen G.Q., Chang C., Bleecker A.B. and Meyerowitz E.M. (1998) *ETR2* is an *ETR1*-like gene involved in ethylene signal transduction in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* 95: 5812–5817.

Sakai, T., Honing H., Nishioka M., Uehara Y., Takahashi M., Fujisawa N., Saji K., Seki M., Shinozaki K., Jones M. A., Smirnoff N., Okada K. and Wasteneys G. O. (2008) Armadillo repeat-containing kinesins and a NIMA-related kinase are required for epidermal-cell morphogenesis in *Arabidopsis*. *Plant J.* 53: 157–171.

Sambrook, J., Fritsch, E.F. and Maniatis, T. (1989). Molecular cloning: a laboratory manual. *Cold Spring Harbor Laboratory Press, Cold Spring Harbor Lab, New York*. 2nd edition.

Schifelbein J. W. and Somerville C. (1990) Genetic Control of Root Hair Development in *Arabidopsis thaliana*. *The Plant Cell* 2(3): 235-243.

Schifelbein J. W., Shipley A. L. and Rowse P. (1992) Calcium influx at the tip of growing root-hair cells of *Arabidopsis thaliana*. *Planta* 4: 455-459.

Schifelbein J., Galway M., Masucci J. and Ford S. (1993) Pollen Tube and Root-Hair Tip Growth Is Disrupted in a Mutant of *Arabidopsis thaliana*. *Plant Physiology* 103(3): 979-985.

Scholl RL, May ST, Ware DH. (2000) Seed and molecular resources for Arabidopsis. *Plant Physiol.* 124(4):1477-80.

Schultz J., Milpetz F., Bork P. and Ponting C. P. (1998) SMART, a simple modular architecture research tool: Identification of signaling domains. *PNAS* 95(11): 5857-5864.

Sexton, R. and Roberts, J. A. (1982). Cell biology of abscission. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 33: 133-162.

Sherrier D. J. and VandenBosch K. A. (1994) Secretion of cell wall polysaccharides in *Vicia* root hairs. *Plant journal* 5(2): 185-195.

Showalter A. M. (2001) Arabinogalactan-proteins: structure, expression and function. *Cellular and Molecular Life Sciences* 58: 1399-1417.

Showalter A. M. (2001) Arabinogalactan-proteins: structure, expression and function. *Cell Mol Life Sci.* 58(10):1399-417. Review.

Sieberer B. J., Timmers A. C. and Emons A. M. (2005) Nod factors alter the microtubule cytoskeleton in *Medicago truncatula* root hairs to allow root hair reorientation. *Mol Plant Microbe Interact* 18(11):1195-204.

Stenvik, G. E., Butenko, M. A., Urbanowicz, B. R., Rose, F. K. C. and Aalen, R. B. (2006). Overexpression of INFLORESCENCE DEFICIENT IN ABSCISSION activates cell separation in vestigial AZs in *Arabidopsis*. *Plant Cell* 18: 1467-1476.

Szymkowiak E. J. and Irish E. E. (2006) Interactions between *jointless* and wild-type tomato tissues during development of the pedicel abscission zone and the inflorescence meristem. *Plant Cell* 11: 159-175.

Szymkowiak E. J. and Irish E. E. (2006) *JOINTLESS* suppresses sympodial identity in inflorescence meristems of tomato. *Planta* 223: 646-658.

Thole J. M., Vermeer J.E., Zhang Y., Gadella T.W. Jr. and Nielsen E. (2008) Root hair defective4 encodes a phosphatidylinositol-4-phosphate phosphatase required for proper root hair development in *Arabidopsis thaliana*. *Plant Cell* 20(2):381-95.

Tucker M. L., Sexton R., del Campillo E. and Lewis L. N. (1988) Bean abscission cellulose. *Plant Physiology* 88(4): 1257-1262.

Visser J. and Voragen A.G.J. (1996) Pectins and Pectinases: Progress in Biotechnology, Vol. 14

Volko SM, Boller T, Ausubel FM. (1998) Isolation of new *Arabidopsis* mutants with enhanced disease susceptibility to *Pseudomonas syringae* by direct screening. *Genetics* 149: 537-48.

Werner S., Beer H. D., Mauch C., Lüscher B. and Werner S. (2001) The Mad1 transcription factor is a novel target of activin and TGF-beta action in keratinocytes: possible role of Mad1 in wound repair and psoriasis. *Oncogene* 20(51):7494-504.

White C.N., Rivin C.J. (1995) Sequence and regulation of a late embryogenesis abundant group 3 protein of maize. *Plant Physiol.* 108: 1337-1338.

Wilson A. K., Pickett F. B., Turner J. C. and Estelle M. (1990) A dominant mutation in *Arabidopsis* confers resistance to auxin, ethylene and abscisic acid. *Molecular and General Genetics* 222(2-3): 377-383.

Wymer C. L., Bibikova T. N. and Gilroy S. (1997) Cytoplasmic free calcium distributions during the development of root hairs of *Arabidopsis thaliana*. *The Plant Journal* 12(2): 427–439.

Yang G., Gao P., Zhang H., Huang S. and Zheng Z. (2007) A Mutation in *MRH2* Kinesin Enhances the Root Hair Tip Growth Defect Caused by Constitutively Activated ROP2 Small GTPase in *Arabidopsis*. *Plos one* 10: e1074.

Zheng Z. L. (2007) Use of the gl1 Mutant & the CA-rop2 Transgenic Plants of *Arabidopsis thaliana* in the Biology Laboratory Course. *Am Biol Teach* 68(9):148-153.

Zhu Z., Shendure J. and Church G. M. (2005) Discovering functional transcription-factor combinations in the human cell cycle. *Cold Spring Harbor Laboratory Press* 15:848–855.

Zimmermann P., Hennig L. and Gruissem W. (2004) Gene-expression analysis and network discovery using Genevestigator. *Trends Plant Sci.* 10(9):407-9.